

# ON $p$ -ADIC CHARACTER DEDEKIND SUMS

Mehmet Cenkci

Communicated by Adnan Tercan

MSC 2010 Classification: 11F20.

Keywords and phrases: Dedekind sums, character Dedekind sums,  $p$ -adic measure theory.

**Abstract** Using  $p$ -adic measure theory we give explicit representations of  $p$ -adic analogues of the character Dedekind sums and their reciprocity laws.

## 1. Introduction

K. Rosen and W. Synder ([12]) showed that by  $p$ -adically interpolating certain partial zeta functions, it is possible to interpolate the higher order Dedekind sums

$$s_m(h, k) = \sum_{a=0}^{k-1} \frac{a}{k} \bar{B}_m \left( \frac{ha}{k} \right)$$

introduced by Apostol ([1]), thus obtain  $p$ -adic Dedekind sums. The authors then showed that there is a reciprocity law for  $p$ -adic Dedekind sums, however they are not able to obtain an explicit form for the reciprocity law for the arbitrary  $p$ -adic integers. C. Synder ([16]) obtained such an explicit form for the reciprocity law for the arbitrary  $p$ -adic integers by the use of  $p$ -adic measure theory. In a series papers A. Kudo ([8, 9, 10]) extended the results of Rosen and Synder to higher order Dedekind sums

$$s_{m+1}^{(r)}(h, k) = \sum_{a=0}^{k-1} \bar{B}_{m+1-r} \left( \frac{a}{k} \right) \bar{B}_r \left( \frac{ha}{k} \right), 0 \leq r \leq m + 1$$

for every  $h, k$  and  $r \geq 1$ . Kudo accomplished this by using an expression for  $k^m s_{m+1}^{(r)}(h, k)$  in terms of Euler numbers and a  $p$ -adic continuous function which interpolates these numbers.

B. Berndt ([2]) gave a character transformation formula similar to those for the Dedekind eta function and defined Dedekind sums with character  $s(h, k; \chi)$  by

$$s(h, k; \chi) = \sum_{a=0}^{kf-1} \chi(a) \bar{B}_1 \left( \frac{a}{kf} \right) \bar{B}_{1,\chi} \left( \frac{ha}{k} \right)$$

for  $(h, k) = 1$ , where  $\chi$  is a primitive Dirichlet character of conductor  $f$  and  $\bar{B}_{m,\chi}(x)$  is the  $m$ th character Bernoulli function. M. Cenkci, M. Can and V. Kurt ([5]) extended this definition as

$$s_m(h, k; \chi) = \sum_{a=0}^{kf-1} \chi(a) \bar{B}_1 \left( \frac{a}{kf} \right) \bar{B}_{m,\chi} \left( \frac{ha}{k} \right) \tag{1.1}$$

and established reciprocity law.

The purpose of this paper is to define  $p$ -adic character Dedekind sums which interpolate (1.1). The basic idea is to use an expression for  $s_m(h, k; \chi)$  in terms of generalized Euler numbers and a  $p$ -adic continuous function which interpolates these numbers. We also show that there is a reciprocity law for these sums for  $m + 1 \equiv 0 \pmod{p - 1}$ , where  $p$  is an odd prime number.

## 2. Preliminaries

For integers  $m, h$  and  $k$  such that  $m \geq 0$  and  $k > 0$  the higher order Dedekind sums are defined as

$$s_m(h, k) = \sum_{a=0}^{k-1} \frac{a}{k} \bar{B}_m \left( \frac{ha}{k} \right), \tag{2.1}$$

where  $\overline{B}_m(x)$  denotes the  $m$ th periodic Bernoulli function defined by

$$\sum_{m=0}^{\infty} B_m(x) \frac{t^m}{m!} = \frac{te^{xt}}{e^t - 1}$$

for all real  $x$  and  $\overline{B}_m(x) = \overline{B}_m(\{x\})$  with  $\{x\}$  denotes the fractional part of  $x$ . For  $x = 0$ ,  $B_m(0) = B_m$  is the  $m$ th Bernoulli number. For even  $m$ , the higher order Dedekind sums are relatively uninteresting. However, for odd  $m$ , they possess a reciprocity formula.

There are other representations of Dedekind sums. Let  $E_m(u)$  be the modified Euler numbers belonging to a parameter  $u$ .  $E_m(u)$  is defined by ([10])

$$\frac{u}{e^t - u} = \sum_{m=0}^{\infty} E_m(u) \frac{t^m}{m!}.$$

Note that  $mE_{m-1}(u) = B_m$  and  $\frac{1-u}{u}E_m(u) = H_m(u)$  for all  $m \in \mathbb{N}$ , where  $H_m(u)$  is the Eulerian number with parameter  $u$  ([4]). It is known that (see [4]) for any  $a \in \mathbb{Z}$ ,  $k, m \in \mathbb{N}$  and for any  $k$ th root of unity  $\zeta$ , we have

$$mE_{m-1}(\zeta) = k^{m-1} \sum_{j=0}^{k-1} \overline{B}_m\left(\frac{j}{k}\right) \zeta^{-j}$$

and

$$k^m \overline{B}_m\left(\frac{a}{k}\right) = m \sum_{\zeta^k=1} E_{m-1}(\zeta) \zeta^{-a}.$$

Now, since  $\frac{a}{k} = \overline{B}_1\left(\frac{a}{k}\right)$ , we have from (2.1) that

$$k^m s_m(h, k) = m \sum_{\zeta^k=1} \frac{E_{m-1}(\zeta)}{\zeta^h - 1}$$

after a little reduction ((6.6) of [3]).

There are many generalizations of Bernoulli numbers and polynomials. One of them is via a Dirichlet character. Let  $\chi$  be a primitive Dirichlet character of conductor  $f$ . Then character Bernoulli polynomials  $B_{m,\chi}(x)$  are defined by

$$\sum_{a=0}^{f-1} \frac{\chi(a) te^{(a+x)t}}{e^{ft} - 1} = \sum_{m=0}^{\infty} B_{m,\chi}(x) \frac{t^m}{m!}.$$

This definition immediately leads the relation

$$B_{m,\chi}(x) = f^{m-1} \sum_{a=0}^{f-1} \chi(a) B_m\left(\frac{a+x}{f}\right).$$

Character Dedekind sums, which we are going to use for the definition of  $p$ -adic character Dedekind sums, are defined by

$$s_m(h, k; \chi) = \sum_{a=0}^{kf-1} \chi(a) \overline{B}_1\left(\frac{a}{kf}\right) \overline{B}_{m,\chi}\left(\frac{ha}{k}\right),$$

where  $\overline{B}_{m,\chi}(x) = B_{m,\chi}(\{x\})$ . We note that for a principal character  $\chi$  this definition reduces to Apostol's.

For a primitive Dirichlet character  $\chi$  of conductor  $f$  let  $E_{m,\chi}(u)$  be the numbers defined by

$$\sum_{a=0}^{f-1} \frac{\chi(a) u^{f-a} e^{at}}{e^{ft} - u^f} = \sum_{m=0}^{\infty} E_{m,\chi}(u) \frac{t^m}{m!}.$$

Note that  $mE_{m-1,\chi}(1) = B_{m,\chi}$  for all  $m \in \mathbb{N}$ . Let  $\zeta$  be an arbitrary primitive  $f$ th root of unity. Then, if  $(k, f) = 1$ , we deduce that

$$\chi(k) k^m \overline{B}_{m,\chi}\left(\frac{a}{k}\right) = m \sum_{\zeta^k=1} E_{m-1,\chi}(\zeta) \zeta^a$$

(Proposition 3.1 of [7]). From this relation we may write the character Dedekind sums in terms of generalized Euler numbers as

$$\chi(k) k^m s_m(h, k; \chi) = \sum_{a=0}^{kf-1} \chi(a) \overline{B}_1\left(\frac{a}{kf}\right) m \sum_{\zeta^k=1} E_{m-1, \chi}(\zeta^h) \zeta^a.$$

Throughout this paper we use standard terminology from the  $p$ -adic theory. Let  $p$  denote a fixed prime number which, for convenience, we assume to be odd. Let  $\mathbb{Z}_p$  and  $\mathbb{Q}_p$  denote the set of  $p$ -adic integers and  $p$ -adic rational numbers, respectively. Let  $|\cdot|_p$  denote the  $p$ -adic absolute value on  $\mathbb{Q}_p$ , normalized so that  $|p|_p = p^{-1}$ . Let  $\overline{\mathbb{Q}}_p$  be the algebraic closure of  $\mathbb{Q}_p$  and  $\mathbb{C}_p$  be the completion of  $\overline{\mathbb{Q}}_p$  with respect to  $p$ -adic absolute value. Note that two fields  $\mathbb{C}$  and  $\mathbb{C}_p$  are algebraically isomorphic, and any one of the two can be embedded in the other.

The group of  $p$ -adic units is denoted by  $\mathbb{Z}_p^*$ . If  $V$  is the group  $\{x \in \mathbb{Q}_p : x^{p-1} = 1\}$ , then  $\mathbb{Z}_p^* = V \times (1 + p\mathbb{Z}_p)$ . Thus, if  $a \in \mathbb{Z}_p^*$  then  $a = w(a) \langle a \rangle$ , where  $w(a)$  and  $\langle a \rangle$  are the projections of  $a$  onto  $V$  and  $1 + p\mathbb{Z}_p$ , respectively. Letting  $w(a) = 0$  for  $a \in \mathbb{Z}$  such that  $(a, p) \neq 1$ , we see that  $w$  is actually a Dirichlet character, called Teichmüller character, having conductor  $p$ . We note that the order of  $w$  is  $p - 1$ .

Let  $f$  be a positive integer. We set  $X_f = \varprojlim_N (\mathbb{Z}/fp^N\mathbb{Z})$ , the map from  $\mathbb{Z}/fp^M\mathbb{Z}$  to  $\mathbb{Z}/fp^N\mathbb{Z}$  for  $M \geq N$ , to be reduction mod  $fp^N$ . In the special case  $f = 1$ ,  $X_1 = \mathbb{Z}_p$ . Let  $a + p^N\mathbb{Z}_p = \{x \in \mathbb{Q}_p : |x - a|_p \leq p^{-N}\}$  for  $a \in \mathbb{Q}_p$  and  $N \in \mathbb{Z}$ . Then the sets of the form  $a + p^N\mathbb{Z}_p$  form a basis of open sets for the metric space  $\mathbb{Q}_p$ . This means that any open subset of  $\mathbb{Q}_p$  is a union of open sets of this type. Note that  $a + fp^N\mathbb{Z}_p = \bigcup_{0 \leq b < p} (a + bfp^N) + fp^{N+1}\mathbb{Z}_p$  and  $X_f^* = X_f \setminus pX_f = \bigcup_{\substack{0 < a < fp \\ (a, p) = 1}} a + fp\mathbb{Z}_p$  (see [6, 13]).

Let  $UD(\mathbb{Z}_p, \mathbb{C}_p)$  be the Banach algebra of all uniformly (or strictly) differentiable functions  $f : \mathbb{Z}_p \rightarrow \mathbb{C}_p$  under the pointwise operations and valuation (see [11, 13, 18]). If  $\zeta$  satisfies the condition that  $\zeta^{p^n} \neq 1$  for all  $n \geq 0$  we can define a finitely additive measure  $\mu_\zeta$  on  $X_f$  by

$$\mu_\zeta(a + fp^N\mathbb{Z}_p) = \frac{\zeta^{fp^N - a}}{1 - \zeta^{fp^N}}, 0 \leq a < fp^N, N \geq 0.$$

Then for a function  $f \in UD(\mathbb{Z}_p, \mathbb{C}_p)$  we have

$$\int_{X_f} f(x) d\mu_\zeta(x) = \lim_{N \rightarrow \infty} \sum_{a=0}^{fp^N-1} f(a) \mu_\zeta(a + fp^N\mathbb{Z}_p) = \lim_{N \rightarrow \infty} \frac{\zeta^{fp^N}}{1 - \zeta^{fp^N}} \sum_{a=0}^{fp^N-1} f(a) \zeta^{-a}$$

using the  $p$ -adic limit of the  $N$ th Riemann sum of  $f$ . Main results for this formula can be given as follows:

**Proposition 2.1.** (see [14, 15, 17]) *Let  $|t|_p \leq p^{1/(1-p)}$ ,  $t \in \mathbb{C}_p$  and  $t \neq 0$ , and let  $\chi$  be a primitive Dirichlet character with conductor  $f$ . Then we have*

- (1)  $E_m(\zeta) = \int_{\mathbb{Z}_p} x^m d\mu_\zeta(x)$ .
- (2)  $E_m(\zeta) - p^m E_m(\zeta^p) = \int_{\mathbb{Z}_p^*} x^m d\mu_\zeta(x)$ .
- (3)  $E_{m, \chi}(\zeta) = \int_{X_f} \chi(x) x^m d\mu_\zeta(x)$ .
- (4)  $E_{m, \chi}(\zeta) - \chi(p) p^m E_{m, \chi}(\zeta^p) = \int_{X_f^*} \chi(x) x^m d\mu_\zeta(x)$ .

### 3. $p$ -adic Interpolation of Character Dedekind Sums and Their Reciprocity Formula

In this section we define a  $p$ -adic continuous function that will be used to interpolate  $s_m(h, k; \chi)$ . Let  $\zeta$  be a root of unity and  $\zeta^{p^n} \neq 1$  for all  $n \geq 0$ . Let

$$F_p(s; \zeta, \chi) = \int_{X_f^*} \chi(x) w^{-1}(x) \langle x \rangle^s \frac{1}{x} d\mu_\zeta(x)$$

for  $s \in \mathbb{Z}_p$ . Let  $\exp$  and  $\log$  denote the  $p$ -adic exponential and logarithm functions respectively. Then, since  $\langle x \rangle \equiv 1 \pmod{p}$  for  $x \in \mathbb{Z}_p^\times$ ,  $\log \langle x \rangle \equiv 0 \pmod{p}$  and  $\langle x \rangle^s = \exp(\log \langle x \rangle)$ . Furthermore, fixing an embedding of algebraic closure of  $\mathbb{Q}, \overline{\mathbb{Q}}$ , into  $\mathbb{C}_p$ , we may then consider the values of a Dirichlet character  $\chi$  as lying in  $\mathbb{C}_p$ . Therefore  $F_p(s; \zeta, \chi)$  is an analytic function of  $s$  in  $\mathbb{Z}_p$  with the expansion

$$F_p(s; \zeta, \chi) = \sum_{m=0}^{\infty} c_m(\zeta, \chi) s^m,$$

$$c_m(\zeta, \chi) = \int_{X_f^*} \chi(x) w^{-1}(x) \frac{(\log \langle x \rangle)^m}{m!} \frac{1}{x} d\mu_\zeta(x),$$

$$|c_m(\zeta, \chi)|_p \leq \left| \frac{p^m}{m!} \right|_p \leq p^{-m} p^{\frac{m}{p-1}}.$$

Now, since the order of  $w$  is  $p - 1$ , we have by (4) of Proposition 2.1 that

$$F_p(m, \zeta, \chi) = \int_{X_f^*} \chi(x) x^{m-1} d\mu_\zeta(x) = E_{m-1, \chi}(\zeta) - \chi(p) p^{m-1} E_{m-1, \chi}(\zeta^p)$$

for all integers  $m \geq 1$  and  $m + 1 \equiv 0 \pmod{p - 1}$ .

**Definition 3.1.** Let  $\chi$  be a Dirichlet character of conductor  $f$  and let  $h \in \mathbb{Z}, k \in \mathbb{N}$  with  $(k, f) = 1$ . Then

$$S_p(s; h, k; \chi) = s \sum_{a=0}^{kf-1} \chi(a) \frac{a}{kf} \sum_{\zeta^k=1} F_p(s; \zeta^h, \chi) \zeta^a$$

is the  $p$ -adic character Dedekind sum for all  $s \in \mathbb{Z}_p$ .

We now show that the function  $S_p(s; h, k; \chi)$  interpolates the character Dedekind sums.

**Proposition 3.2.** Let  $\chi$  be a Dirichlet character of conductor  $f$ . For any integers  $m, h, k$  such that  $m \geq 0, m + 1 \equiv 0 \pmod{p - 1}, k > 0$  and  $(k, f) = 1$  we have

$$S_p(m; h, k; \chi) = \chi(k) k^m \{ s_m(h, k; \chi) - \chi(p) p^{m-1} s_m(ph, k; \chi) \}.$$

*Proof.* Proof follows from definitions of  $S_p(s; h, k; \chi)$  and  $F_p(s; \zeta, \chi)$ . In fact we have

$$\begin{aligned} S_p(m; h, k; \chi) &= \sum_{a=0}^{kf-1} \chi(a) \frac{a}{kf} m \sum_{\zeta^k=1} F_p(m; \zeta^h, \chi) \zeta^a \\ &= \sum_{a=0}^{kf-1} \chi(a) \frac{a}{kf} m \sum_{\zeta^k=1} [E_{m-1, \chi}(\zeta^h) - \chi(p) p^{m-1} E_{m-1, \chi}(\zeta^{hp})] \zeta^a \\ &= \sum_{a=0}^{kf-1} \chi(a) \frac{a}{kf} m \sum_{\zeta^k=1} E_{m-1, \chi}(\zeta^h) \zeta^a \\ &\quad - \chi(p) p^{m-1} \sum_{a=0}^{kf-1} \chi(a) \frac{a}{kf} m \sum_{\zeta^k=1} E_{m-1, \chi}(\zeta^{hp}) \zeta^a \\ &= \chi(k) k^m s_m(h, k; \chi) - \chi(k) k^m \chi(p) p^{m-1} s_m(ph, k; \chi), \end{aligned}$$

which is the result. □

Now we are going to interpolate the reciprocity law for  $s_m(h, k; \chi)$ . For odd integer  $m$ , coprime positive integers  $h$  and  $k$ , a non-principle primitive Dirichlet character  $\chi$  of modulus  $f$  and  $(f, hk) = 1$  we have the reciprocity formula (see [5])

$$\begin{aligned} &hk^m s_m(h, k; \chi) + kh^m s_m(k, h; \bar{\chi}) \\ &= \frac{1}{m+1} \sum_{j=0}^{m+1} \binom{m+1}{j} h^j k^{m+1-j} B_{j, \bar{\chi}} B_{m+1-j, \chi} + \frac{m}{f} \chi(k) \bar{\chi}(-h) (f^{m+1} - 1) B_{m+1}. \end{aligned}$$

**Proposition 3.3.** *Let  $m$  be an odd integer,  $h$  and  $k$  coprime positive integers,  $\chi$  be a non-principle primitive Dirichlet character of modulus  $f$ ,  $k \equiv 1 \pmod{f}$ ,  $h \equiv 1 \pmod{f}$  and  $p$  be an odd prime number with  $(p, kf) = (p, hf) = 1$ ,  $m + 1 \equiv 0 \pmod{p-1}$ . Then we have*

$$hS_p(m; h, k; \chi) + kS_p(m; k, h; \bar{\chi}) = (1 - p^{m-1}) \{hk^m s_m(h, k; \chi) + kh^m s_m(k, h; \bar{\chi})\}.$$

*Proof.* From Proposition 3.2 we have

$$\begin{aligned} S_p(m; h, k; \chi) &= \chi(k) k^m \{s_m(h, k; \chi) - \chi(p) p^{m-1} s_m(ph, k; \chi)\}, \\ S_p(m; k, h; \bar{\chi}) &= \bar{\chi}(h) h^m \{s_m(k, h; \bar{\chi}) - \bar{\chi}(p) p^{m-1} s_m(pk, h; \bar{\chi})\}. \end{aligned}$$

Since  $k \equiv 1 \pmod{f}$  and  $h \equiv 1 \pmod{f}$  we have  $\chi(k) = \bar{\chi}(h) = 1$ . Thus

$$\begin{aligned} hS_p(m; h, k; \chi) + kS_p(m; k, h; \bar{\chi}) &= hk^m s_m(h, k; \chi) + kh^m s_m(k, h; \bar{\chi}) \\ &\quad - p^{m-1} \{\chi(p) hk^m s_m(ph, k; \chi) + \bar{\chi}(p) kh^m s_m(pk, h; \bar{\chi})\}. \end{aligned}$$

Now

$$\begin{aligned} \chi(p) s_m(ph, k; \chi) &= \chi(p) \sum_{a=1}^{kf} \chi(a) \bar{B}_1\left(\frac{a}{kf}\right) \bar{B}_{m, \chi}\left(\frac{pha}{k}\right) \\ &= \sum_{a=1}^{kf} \chi(pa) \bar{B}_1\left(\frac{a}{kf}\right) \bar{B}_{m, \chi}\left(\frac{hpa}{k}\right). \end{aligned}$$

Since  $(p, kf) = 1$ , we may write  $\bar{B}_1\left(\frac{a}{kf}\right) = \bar{B}_1\left(\frac{pa}{kf}\right)$ , and the values  $pa$  run through the same values of  $a$ . Therefore we have

$$\chi(p) s_m(ph, k; \chi) = s_m(h, k; \chi).$$

Similarly

$$\bar{\chi}(p) s_m(pk, h; \bar{\chi}) = s_m(k, h; \bar{\chi}),$$

which completes the proof.  $\square$

## References

- [1] T. M. Apostol, *Generalized Dedekind sums and transformation formulae of certain Lambert series*, Duke Math. J. **17**, 147–157 (1950).
- [2] B. C. Berndt, *Character transformation formulae similar to those for the Dedekind Eta-function*, in ‘Analytic Number Theory’, Proc. Sym. Pure Math. XXIV, Amer. Math. Soc., Providence R. I., 9–30 (1973).
- [3] L. Carlitz, *Some theorems on generalized Dedekind sums*, Pacific J. Math. **3**, 513–522 (1953).
- [4] L. Carlitz, *Eulerian numbers and polynomials*, Math. Mag. **32**, 247–260 (1959).
- [5] M. Cenkci, M. Can, V. Kurt, *Degenerate and character Dedekind sums*, J. Number Theory **124**, 346–363 (2007).
- [6] N. Koblitz,  *$p$ -adic Numbers,  $p$ -adic Analysis and Zeta Functions*, Graduate Texts in Mathematics, 58, Springer-Verlag, Berlin, Heidelberg, New York, (1984).
- [7] K. Kozuka, *Dedekind type sums attached to Dirichlet characters*, Kyushu J. Math. **58**, 1–24 (2004).
- [8] A. Kudo, *On  $p$ -adic Dedekind sums (II)*, Mem. Fac. Sci. Kyushu Univ. **45**, 245–284 (1991).
- [9] A. Kudo, *Reciprocity formulas for  $p$ -adic Dedekind sums*, Bull. Fac. Lib. Arts Nagasaki Univ. Nat. Sci. **34**, 97–101 (1994).
- [10] A. Kudo, *On  $p$ -adic Dedekind sums*, Nagoya Math. J. **144**, 155–170 (1996).
- [11] A. M. Robert, *A Course in  $p$ -Adic Analysis*, Springer-Verlag, New York, (2000).
- [12] K. Rosen, W. Synder,  *$p$ -adic Dedekind sums*, J. Reine Angew. Math. **361**, 23–26 (1985).
- [13] W. H. Schikhof, *Ultrametric Calculus, An introduction to  $p$ -adic analysis*, Cambridge Studies in Adv. Math. 4, Cambridge University Press, Cambridge, New York, (1984).
- [14] K. Shiratani, *On Euler numbers*, Mem. Fac. Sci. Kyushu Univ. **27**, 1–5 (1973).
- [15] K. Shiratani, S. Yamamoto, *On a  $p$ -adic interpolation function for the Euler numbers and its derivatives*, Mem. Fac. Sci. Kyushu Univ. **39**, 113–125 (1985).
- [16] C. Synder,  *$p$ -adic interpolation of Dedekind sums*, Bull. Austral. Math. Soc. **38**, 293–301 (1988).

- 
- [17] H. Tsumura, *On a  $p$ -adic interpolation of the generalized Euler numbers and its applications*, Tokyo J. Math. **10**, 281–293 (1987).
- [18] C. F. Woodcock, *An invariant  $p$ -adic integral on  $\mathbb{Z}_p$* , J. London Math. Soc., **8**, 731–734 (1974).

**Author information**

Mehmet Cenkci, Akdeniz University, Department of Mathematics, Antalya, 07058, Turkey.  
E-mail: [cenkci@akdeniz.edu.tr](mailto:cenkci@akdeniz.edu.tr)

Received: March 13, 2015.

Accepted: May 20, 2015.