

# New Type of Difference Triple Sequence Spaces

Shyamal Debnath and Bimal Chandra Das

Communicated by Faud Kittaneh

MSC 2010 Classifications: 40A05, 46A45.

Keywords and phrases: triple sequence space, difference operator, solidity, symmetricity.

The authors are highly grateful to the anonymous referee for his/her valuable comments and suggestions for the improvement of the paper.

**Abstract.** In this article we have introduced the difference triple sequence spaces  $c_0^3(\Delta_p)$ ,  $c^3(\Delta_p)$ ,  $c^{3R}(\Delta_p)$ ,  $l_\infty^3(\Delta_p)$  and  $c^{3B}(\Delta_p)$ , applying the difference operator  $\Delta_p$  for a fixed  $p \in \mathbf{N}$  on the sequence  $(x_{lmn})$  and examine whether the spaces being symmetric, solid, convergence free etc. We have also obtained and proved some inclusion relation too.

## 1. Introduction and Preliminaries :

A triple sequence (real or complex) can be defined as a function  $x : N \times N \times N \rightarrow R(C)$ , where  $N, R$  and  $C$  denote the set of natural numbers, real numbers and complex numbers respectively. The different types of notions of triple sequences was introduced and investigated at the initial stage by Sahiner et. al. [2] and Dutta et. al. [4] and others. Recently Savas and Esi [7] have introduced statistical convergence of triple sequences on probabilistic normed space. Later on, Esi [1] have introduced statistical convergence of triple sequences in topological groups.

The notion of difference sequence spaces (for single sequences) was introduced by Kizmaz [8] as follows:

$Z(\Delta) = \{(x_n) \in w : (\Delta x_n) \in Z\}$ , for  $Z = c, c_0, l_\infty$ , the spaces of convergent, null and bounded sequences, respectively, where  $\Delta x_n = x_n - x_{n+1}$  for all  $n \in N$ . Tripathy and Esi[5] introduced the notion of difference sequence space as  $\Delta_m x = (\Delta_m x_n) = x_n - x_{n+m}$  for all  $n \in \mathbf{N}$  and  $m \in \mathbf{N}$  be fixed. Later on it was studied by Tripathy and Sarma [6] and introduced difference double sequence spaces as follows: .

$Z(\Delta) = \{(x_{mn}) \in w : (\Delta x_{mn}) \in Z\}$ , for  $Z = c^2, c_0^2, l_\infty^2$ , the spaces of convergent, null and bounded double sequences respectively, where  $\Delta x_{mn} = x_{mn} - x_{mn+1} - x_{m+1n} + x_{m+1n+1}$  for all  $m, n \in \mathbf{N}$ .

**Definition 1.1** [2]: A triple sequence  $(x_{lmn})$  is said to be convergent to  $L$  in Pringsheim's sense if for every  $\epsilon > 0$ , there exists  $N(\epsilon) \in N$  such that .

$$|x_{lmn} - L| < \epsilon \text{ whenever } l \geq N, m \geq N, n \geq N \text{ and we write } \lim_{l,m,n \rightarrow \infty} x_{lmn} = L.$$

**Note:** A triple sequence is convergent in Pringsheim's sense may not be bounded [2].

**Definition 1.2** [2]: A triple sequence  $(x_{lmn})$  is said to be Cauchy sequence if for every  $\epsilon > 0$ , there exists  $N(\epsilon) \in N$  such that.

$$|x_{lmn} - x_{pqr}| < \epsilon \text{ whenever } l \geq p \geq N, m \geq q \geq N, n \geq r \geq N .$$

**Definition 1.3** [2]: A triple sequence  $(x_{lmn})$  is said to be bounded if there exists  $M > 0$ , such that  $|x_{lmn}| < M$  for all  $l, m, n \in N$ .

**Definition 1.4** [4]: A triple sequence  $(x_{lmn})$  is said to be converge regularly if it is convergent in Pringsheim's sense and in addition the following limits holds:

$$\lim_{n \rightarrow \infty} x_{lmn} = L_{lm} \quad (l, m \in \mathbf{N})$$

$$\lim_{m \rightarrow \infty} x_{lmn} = L_{ln} \quad (l, n \in \mathbf{N})$$

$$\lim_{l \rightarrow \infty} x_{lmn} = L_{mn} \quad (m, n \in \mathbf{N})$$

Let  $w^3$  denote the set of all triple sequence of real numbers. Then the class of triple sequences  $c_0^3, c^3, l_\infty^3, c^{3R}$  and  $c^{3B}$  denotes the triple sequence spaces which are convergent to zero in Pringsheim's sense, convergent in Pringsheim's sense, bounded in Pringsheim's sense, regularly convergent, bounded and convergent respectively.

These classes are all linear spaces.

It is obvious that  $c_0^3 \subset c^3, c^{3R} \subset c^{3B} \subset l_\infty^3$  and the inclusion are strict.

**Theorem 1.1:** The spaces  $c_0^3, c^3, l_\infty^3, c^{3R}$  and  $c^{3B}$  are complete normed linear spaces with the normed.

$$\|x\| = \sup_{l,m,n} |x_{lmn}| < \infty$$

**Proof:** simple.

**Example 1.1 [2]:** Let  $x_{lmn} = \begin{cases} mn, & l = 3 \\ nl, & m = 5 \\ lm, & n = 7 \\ 8, & \text{otherwise} \end{cases}$

Then  $(x_{lmn}) \rightarrow 8$  in Pringsheim's sense but not bounded as well as not regularly convergent.

**Example 1.2:** Let  $x_{lmn} = 1$ , for all  $l, m, n \in \mathbf{N}$ . Then  $(x_{lmn})$  is convergent in Pringsheim's sense, bounded and regularly convergent.

**Definition 1.5 [4]:** A triple sequence space  $E$  is said to be solid if  $(\alpha_{lmn}x_{lmn}) \in E$  whenever  $(x_{lmn}) \in E$  and for all sequences  $(\alpha_{lmn})$  of scalars with  $|\alpha_{lmn}| \leq 1$ , for all  $l, m, n \in \mathbf{N}$ .

**Definition 1.6 [4]:** A triple sequence space  $E$  is said to be monotone if it contains the canonical pre-images of all its step spaces.

**Remark 1.1 [4]:** A sequence space is solid implies that it is monotone.

**Definition 1.7 [4]:** A triple sequence space  $E$  is said to be convergence free if  $(y_{lmn}) \in E$ , whenever  $(x_{lmn}) \in E$  and  $x_{lmn} = 0$  implies  $y_{lmn} = 0$ .

**Definition 1.8 [4]:** A triple sequence space  $E$  is said to be symmetric if  $(x_{lmn}) \in E$  implies  $(x_{\pi(l)\pi(m)\pi(n)}) \in E$ , where  $\pi$  is a permutation of  $\mathbf{N} \times \mathbf{N} \times \mathbf{N}$ .

Now we introduced the  $p^{th}$  order difference triple sequence spaces as follows:

$$c_0^3(\Delta_p) = \{ (x_{lmn}) \in w^3 : (\Delta_p x_{lmn}) \text{ is regularly null} \}$$

$$c^3(\Delta_p) = \{ (x_{lmn}) \in w^3 : (\Delta_p x_{lmn}) \text{ is convergent in Pringsheim's sense} \}$$

$$c^{3R}(\Delta_p) = \{ (x_{lmn}) \in w^3 : (\Delta_p x_{lmn}) \text{ is regularly convergent} \}$$

$$l_\infty^3(\Delta_p) = \{ (x_{lmn}) \in w^3 : (\Delta_p x_{lmn}) \text{ is bounded} \}$$

$$c^{3B}(\Delta_p) = \{ (x_{lmn}) \in w^3 : (\Delta_p x_{lmn}) \text{ is convergent in Pringsheim's sense and bounded} \}$$

Where  $\Delta_p x_{lmn} = x_{lmn} - x_{lmn+p} - x_{lm+pn} + x_{lm+pn+p} - x_{l+pmn} + x_{l+pmn+p} + x_{l+pm+pn} - x_{l+pm+pn+p}$

when  $p = 1$ , the above spaces becomes  $c_0^3(\Delta), c^3(\Delta), c^{3R}(\Delta), l_\infty^3(\Delta)$  and  $c^{3B}(\Delta)$ , which was

studied by Debnath, Sarma and Das [10].

**2. Main Result:**

**Theorem 2.1:** The classes of sequences  $c_0^3(\Delta_p)$ ,  $c^3(\Delta_p)$ ,  $c^{3R}(\Delta_p)$ ,  $l_\infty^3(\Delta_p)$ ,  $c^{3B}(\Delta_p)$  are linear spaces.

**Proof:** Obvious.

**Theorem 2.2:** The classes of sequences  $c_0^3(\Delta_p)$ ,  $c^3(\Delta_p)$ ,  $c^{3R}(\Delta_p)$ ,  $l_\infty^3(\Delta_p)$  and  $c^{3B}(\Delta_p)$  are complete normed linear spaces with the norm

$$\|x\| = \sup_l |x_{lpp}| + \sup_m |x_{pmp}| + \sup_n |x_{ppn}| + \sup_{l,m,n} |\Delta_p x_{lmn}| < \infty$$

**Proof:** Let  $(x^i)$  be a Cauchy sequence in  $l_\infty^3(\Delta_p)$ , where  $x^i = (x_{lmn}^i) \in l_\infty^3(\Delta_p)$  for each  $i \in \mathbf{N}$ .

Then we have,

$$\|x^i - x^j\| = \sup_l |x_{lpp}^i - x_{lpp}^j| + \sup_m |x_{pmp}^i - x_{pmp}^j| + \sup_n |x_{ppn}^i - x_{ppn}^j| + \sup_{l,m,n} |\Delta_p x_{lmn}^i - \Delta_p x_{lmn}^j| \rightarrow 0$$

as  $i, j \rightarrow \infty$

Therefore,  $|x_{lmn}^i - x_{lmn}^j| \rightarrow 0$ , for  $i, j \rightarrow \infty$  and each  $l, m, n \in \mathbf{N}$

Hence  $(x_{lmn}^i) = (x_{lmn}^1, x_{lmn}^2, x_{lmn}^3, \dots \dots \dots)$  is a Cauchy sequence in  $\mathbf{R}$  (Real numbers).

Whence by the completeness of  $\mathbf{R}$ , it converges to  $x_{lmn}$  say, i.e., there exists

$$\lim x_{lmn}^i = x_{lmn} \text{ for each } l, m, n \in \mathbf{N}$$

Further for each  $\epsilon > 0$ , there exists  $\mathbf{N} = \mathbf{N}(\epsilon)$ , such that for all  $i, j \geq \mathbf{N}$ , and for all  $l, m, n \in \mathbf{N}$

$$|x_{lpp}^i - x_{lpp}^j| < \epsilon, |x_{pmp}^i - x_{pmp}^j| < \epsilon, |x_{ppn}^i - x_{ppn}^j| < \epsilon$$

$$|\Delta_p x_{lmn}^i - \Delta_p x_{lmn}^j| = |(x_{l+pm+pn+p}^i - x_{l+pm+pn+p}^j) - (x_{l+pm+pn}^i - x_{l+pm+pn}^j) - (x_{l+pmn+p}^i - x_{l+pmn+p}^j) + (x_{l+pmn}^i - x_{l+pmn}^j) - (x_{lm+pn+p}^i - x_{lm+pn+p}^j) + (x_{lm+pn}^i - x_{lm+pn}^j) + (x_{lmn+p}^i - x_{lmn+p}^j) - (x_{lmn}^i - x_{lmn}^j)| < \epsilon$$

and

$$\lim_j |x_{lpp}^i - x_{lpp}^j| = |x_{lpp}^i - x_{lpp}| \leq \epsilon,$$

$$\lim_j |x_{pmp}^i - x_{pmp}^j| = |x_{pmp}^i - x_{pmp}| \leq \epsilon,$$

$$\lim_j |x_{ppn}^i - x_{ppn}^j| = |x_{ppn}^i - x_{ppn}| \leq \epsilon,$$

Now

$$\lim_j |\Delta x_{lmn}^i - \Delta x_{lmn}^j| = |(x_{l+pm+pn+p}^i - x_{l+pm+pn+p}^j) - (x_{l+pm+pn}^i - x_{l+pm+pn}^j) - (x_{l+pmn+p}^i - x_{l+pmn+p}^j) + (x_{l+pmn}^i - x_{l+pmn}^j) - (x_{lm+pn+p}^i - x_{lm+pn+p}^j) + (x_{lm+pn}^i - x_{lm+pn}^j) + (x_{lmn+p}^i - x_{lmn+p}^j) - (x_{lmn}^i - x_{lmn}^j)| \leq \epsilon, \text{ for all } i \geq \mathbf{N}$$

Since  $\epsilon$  is not dependent on  $l, m, n$

$$\sup_{l,m,n} |(x_{l+pm+pn+p}^i - x_{l+pm+pn+p}^j) - (x_{l+pm+pn}^i - x_{l+pm+pn}^j) - (x_{l+pmn+p}^i - x_{l+pmn+p}^j) + (x_{l+pmn}^i - x_{l+pmn}^j) - (x_{lm+pn+p}^i - x_{lm+pn+p}^j) + (x_{lm+pn}^i - x_{lm+pn}^j) + (x_{lmn+p}^i - x_{lmn+p}^j) - (x_{lmn}^i - x_{lmn}^j)| \leq \epsilon,$$

Consequently we have,  $\|x_{lmn}^i - x_{lmn}\| \leq 4\epsilon$ , for all  $i \geq \mathbf{N}$

Hence we obtain  $x_{lmn}^i \rightarrow x_{lmn}$  as  $i \rightarrow \infty$  in  $l_\infty^3(\Delta)$

Now we have to show that  $(x_{lmn}) \in l_\infty^3(\Delta)$

$$|x_{lmn} - x_{l+pm+pn+p}| = |x_{lmn} - x_{lmn}^N + x_{lmn}^N - x_{l+pm+pn+p}^N + x_{l+pm+pn+p}^N - x_{l+pm+pn+p}|$$

$$\leq |x_{lmn}^N - x_{l+pm+pn+p}^N| + \|x_{lmn}^N - x_{lmn}\| = O(1)$$

This implies  $x = (x_{lmn}) \in l_\infty^3(\Delta)$

Since  $l_\infty^3(\Delta)$  is a linear space.

Hence  $l_\infty^3(\Delta)$  is complete.

Similarly the others.

**Theorem 2.3:**

- (i)  $c_0^3(\Delta_p) \subset c^3(\Delta_p)$  and the inclusion is strict. .
- (ii)  $c^{3R}(\Delta_p) \subset c^3(\Delta_p)$  and the inclusion is strict.
- (iii)  $c^{3R}(\Delta_p) \subset c^{3B}(\Delta_p)$  and the inclusion is strict.

**Proof:** The inclusion being strict follows from the following example:

**Example 2.1:** We consider the sequence  $(x_{lmn})$  defined by

$$(x_{lmn}) = lmn, \text{ for all } l, m, n \in \mathbf{N}$$

Then  $x_{lmn} \in c^3(\Delta_p)$ , but  $x_{lmn} \notin c_0^3(\Delta_p)$

Hence the inclusion is strict.

**Example 2.2:** We consider the sequence defined by

$$x_{lmn} = \begin{cases} (-1)^n lmn, & \text{for } l = 1, m = 1, 2, 3, 4 \text{ for all } n \in \mathbf{N} \\ -2, & \text{otherwise} \end{cases}$$

Clearly  $x_{lmn} \in c^3(\Delta_p)$ , but the sequence  $x_{lmn} \notin c^{3R}(\Delta_p)$

Hence the inclusion  $c^{3R}(\Delta_p) \subset c^3(\Delta_p)$ , is strict.

**Example 2.3:** We consider the sequence defined by

$$x_{lmn} = \begin{cases} 2, & \text{when } l = 1, m + n \text{ is even, for all } l, m, n \in \mathbf{N} \\ -1, & \text{when } l = 1, m + n \text{ is odd, for all } l, m, n \in \mathbf{N} \\ 1, & \text{otherwise} \end{cases}$$

$x_{lmn} \in c^{3B}(\Delta_p)$ , but the sequence  $x_{lmn} \notin c^{3R}(\Delta_p)$

Therefore, the inclusion is strict.

**Theorem 2.4:** The classes of sequences  $c_0^3(\Delta_p)$ ,  $c^3(\Delta_p)$ ,  $c^{3R}(\Delta_p)$ ,  $l_\infty^3(\Delta_p)$  and  $c^{3B}(\Delta_p)$  are not solid for  $p$  odd.

**Proof:** This is clear from the following examples:

**Example 2.4:** We consider the sequence  $(x_{lmn})$  defined by

$$(x_{lmn}) = 3, \text{ for all } l, m, n \in \mathbf{N}$$

Clearly the triple sequence  $x_{lmn} \in c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p)$  and  $c^{3B}(\Delta_p)$

Consider the sequence of scalars defined by

$$\alpha_{lmn} = (-1)^{l+m+n}, \text{ for all } l, m, n \in \mathbf{N}$$

Then the sequence  $(\alpha_{lmn}x_{lmn})$  takes the following form

$$\alpha_{lmn}x_{lmn} = 3 \cdot (-1)^{l+m+n}, \text{ for all } l, m, n \in \mathbf{N}$$

Clearly  $(\alpha_{lmn}x_{lmn}) \notin c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p)$  and  $c^{3B}(\Delta_p)$  (for  $p$  odd)

Hence  $c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p)$  and  $c^{3B}(\Delta_p)$  are not solid.

**Example 2.5:** We consider the sequence  $(x_{lmn})$  defined by

$$(x_{lmn}) = lmn, \text{ for all } l, m, n \in \mathbf{N}$$

Clearly the sequence  $x_{lmn} \in l_\infty^3(\Delta_p)$

Consider the sequence of scalars defined by

$$\alpha_{lmn} = (-1)^{l+m+n}, \text{ for all } l, m, n \in \mathbf{N}$$

Then the sequence  $(\alpha_{lmn}x_{lmn})$  takes the following form

$$\alpha_{lmn}x_{lmn} = (-1)^{l+m+n}lmn, \text{ for all } l, m, n \in \mathbf{N}$$

Clearly, when  $p$  odd,  $(\alpha_{lmn}x_{lmn}) \notin l_\infty^3(\Delta_p)$ ,

Hence  $l_\infty^3(\Delta_p)$  are not solid.

**Theorem 2.5:** The spaces  $c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p)$  and  $c^{3B}(\Delta_p)$  are not monotone for  $p$  odd.

**Proof:** The proof is clear from the following example:

**Example 2.6:** Consider the sequence  $(x_{lmn})$  defined by

$$(x_{lmn}) = 1, \text{ for all } l, m, n \in \mathbf{N}$$

It is clear that  $x_{lmn} \in c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p)$  and  $c^{3B}(\Delta_p)$

Now we consider the sequence  $(y_{lmn})$  in its pre-image space defined by

$$y_{lmn} = \begin{cases} 1, & \text{for all } l + m + n \text{ is odd, for all } l, m, n \in \mathbf{N} \\ 0, & \text{otherwise} \end{cases}$$

Clearly when  $p$  odd,  $(y_{lmn}) \notin c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p)$  and  $c^{3B}(\Delta_p)$

Hence the spaces  $c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p)$  and  $c^{3B}(\Delta_p)$  are not monotone.

**Theorem 2.6:** The classes of sequences  $c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p)$  and  $c^{3B}(\Delta_p)$  are not symmetric when  $p$  odd.

**Proof:** The proof is followed by the following example:

**Example 2.7:** Consider the triple sequence  $(x_{lmn})$  defined by

$$(x_{lmn}) = m, \text{ for all } m \in \mathbf{N}$$

Clearly the sequence  $x_{lmn} \in c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p)$  and  $c^{3B}(\Delta_p)$

Consider a rearrange sequence  $(y_{lmn})$  of  $(x_{lmn})$  defined by

$$y_{lmn} = \begin{cases} m + 1, & \text{for } m = l, n \text{ is even} \\ m - 1, & \text{for } m = l + 1, n \text{ is even} \\ m, & \text{otherwise} \end{cases}$$

Clearly  $(y_{lmn}) \notin c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p)$  and  $c^{3B}(\Delta_p)$  (for  $p$  odd)

Hence  $c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p)$  and  $c^{3B}(\Delta_p)$  are not symmetric.

**Theorem 2.7:** When  $p$  is odd, the classes of sequences  $c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p), l_\infty^3(\Delta_p)$  and  $c^{3B}(\Delta_p)$  are not convergent free.

**Proof:** We provide an example to prove the result.

**Example 2.8:** Consider the sequence defined by

$$x_{lmn} = \begin{cases} 0, & \text{if } n = 1, \text{ for all } l, m \in \mathbf{N} \\ 2, & \text{otherwise} \end{cases}$$

Clearly the triple sequence  $x_{lmn} \in c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p), l_\infty^3(\Delta_p)$  and  $c^{3B}(\Delta_p)$

Let the sequence  $(y_{lmn})$  be defined by

$$y_{lmn} = \begin{cases} 0, & \text{if } n \text{ is odd, for all } l, m \in \mathbf{N} \\ lmn, & \text{otherwise} \end{cases}$$

Clearly  $(y_{lmn}) \notin c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p), l_\infty^3(\Delta_p)$  and  $c^{3B}(\Delta_p)$ , (for  $p$  odd)

Hence  $c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p), l_\infty^3(\Delta_p)$  and  $c^{3B}(\Delta_p)$ , are not convergence free.

**Theorem 2.8:** The classes of sequences  $c_0^3(\Delta_p), c^3(\Delta_p), c^{3R}(\Delta_p), l_\infty^3(\Delta_p)$  and  $c^{3B}(\Delta_p)$  are all sequence algebra.

**Proof:** It is obvious.

## References

- [1] A. Esi, Statistical convergence of triple sequences in topological groups, Annals of the University of Craiova, Mathematics and Computer Science Series, 40(1)(2013), 29-33.
- [2] A. Sahiner, M. Gurdal and K. Duden, Triple sequences and their statistical convergence, Selcuk. J. Appl. Math., 8(2) (2007),49-55.
- [3] A. Sahiner, B.C.Tripathy, Some I-related properties of Triple sequences, Selcuk. J. Appl. Math., 9(2) (2008),9-18.
- [4] A.J.Datta, A.Esi, B.C.Tripathy, Statistically convergent triple sequence spaces defined by Orlicz function, J. Math. Anal., 4(2) (2013) 16-22.
- [5] B. C. Tripathy and A. Esi, A new type of difference sequence spaces, International J. Sci. Tech. 1:1(2006),11-14.
- [6] B.C.Tripathy, B. Sarma, Statistically convergent difference double sequence spaces, Acta Math. Sinica, 24(5) (2008),737-742.
- [7] E.Savas and A. Esi, Statistical convergence of triple sequences on probabilistic normed space, Annals of the University of Craiova, Mathematics and Computer Science Series, 39(2)(2012), 226-236.
- [8] H.Kizmaz, On certain sequence spaces, Canad. Math. Bull.,24:2(1981),169-176.

- [9] M. Et and R. Colak, On some generalized difference sequence spaces, *Soochow J. of Math.*, 21(1995), 377-386.
- [10] S. Debnath, B. Sarma and B.C.Das, Some Generalized Triple Sequence Spaces of Real Numbers, accepted, *J. Non linear Anal. Opti.*
- [11] S. Debnath, B. Sarma and S. Saha, Some Sequence Spaces of Interval Vectors, *Afrika Matematika* ( Accepted, online, 2014).
- [12] S. Debnath and J. Debnath, Some generalized statistical convergent sequence spaces of fuzzy numbers via ideals, *Math. Sci. Lett.*, V-2(2), (2013), 151-154.
- [13] S. Debnath and J. Debnath, Some ideal convergent sequence spaces of fuzzy real numbers, *Palestine J. Mathematics*, Vol. 3(1) (2014), 27-32.
- [14] S. Debnath and S. Saha, Some Newly Defined Sequence Spaces Using Regular Matrix Of Fibonacci Numbers, *AKU-J. Sci. Eng.* (Accepted, online, 2014).
- [15] T. Salat, On Statistically Convergent Sequences of Real Numbers, *Math. Slovaca*, 30(1980), No. 2, 139-150.

**Author information**

Shyamal Debnath, Department of Mathematics, Tripura University (A Central University), Suryamaninagar-799022, Agartala, india.

E-mail: shyamalnitamath@gmail.com

Bimal Chandra Das, Department of Mathematics, Govt. Degree College, Kamalpur, Dhalai, Tripura, India.

E-mail: bcdas3744@gmail.com

Received: July 10, 2014.

Accepted: September 23, 2014.