

GRAPH OPERATIONS ON THE SYMMETRIC DIVISION DEG INDEX OF GRAPHS

C. K. Gupta¹, V. Lokesh^{2*}, Shwetha B. S³ and Ranjini P. S⁴

Communicated by P. K. Banerji

MSC 2010 Classifications: Primary: 05C20; Secondary: 05C05.

Keywords and phrases: Graph operations, Symmetric division deg index, Maximum degree, nilpotent matrix group, Cayley graph.

The second and third authors are visiting researchers of Prof. C. K. Gupta, and they thank the authorities of University of Manitoba, Canada for providing an necessary arrangements to complete this work.

Abstract. The Symmetric division deg index of a connected graph G , is defined as $SDD(G) = \sum_{uv \in E(G)} \frac{d_u}{d_v} + \frac{d_v}{d_u}$ where d_v is the degree of a vertex v in G . In this paper, we concentrated on the graph operations like lexicographic product, symmetric difference and corona product of graphs related to the symmetric division deg index.

1 Introduction and Preliminaries

Molecular descriptors, being numerical functions of molecular structure, play a fundamental role mathematical chemistry. They are used in QSAR and QSPR studies to relate biological or chemical properties of molecules to specific molecular descriptors [3]. Topological indices, being numerical functions of the underlying molecular graph, represent an important type of molecular descriptors. Some applications related topological indices related to smart polymers are found in [11] and comparative study of topological indices ad molecular weigh of some carbohydrates are in [12]. Recently in [5], C. K. Gupta and et al., established the relations on graph operations on matrix group.

Inspired by the most successful indices of this form, such as second zagreb index,[4], Randic index [10], [15] and others, there was defined a whole family of Adriatic indices [17]. In recent times [16], D. Vukicevic revealed the set of 148 discrete Adriatic indices. They were analyzed on the testing sets provided by the International Academy of Mathematical Chemistry and it had been shown that they have good predictive properties in many cases. There was a vast research regarding various properties of this topological index.

In a new article [17], D. Vukicevic posed the open questions in his end of the paper. Stimulate from this, here we ardent one of the index specifically, SDD. This emphasizes the development of lower and upper bounds for graphs [6]. This acquires some results that are partial answer to the open queries.

Symmetric division deg index is one of the discrete Adriatic indices that is good predictor of total surface area for polychlorobiphenyls. Some results on symmetric division deg index is also found in [1].

In group theory, a nilpotent group is a group that is "almost abelian". This idea is motivated by the fact that nilpotent groups are solvable and for finite nilpotent groups, two elements having relatively prime orders must commute. The multiplicative group of upper unitriangular $n \times n$ matrices over any field F is a nilpotent group of length $n - 1$ [14]. Here, we discussed some relations related to cayley graph of nilpotent matrix group of length one related to SDD.

We recall some definitions which are essential.

Definition: 1 The first Zagreb index [9] defined as,

$$M_1(G) = \sum d(u)^2 = \sum_{u,v \in E(G)} [d(u) + d(v)].$$

Definition: 2 The Symmetric division deg index of a connected graph G , is defined as

$$SDD(G) = \sum_{uv \in E(G)} \frac{\max(d_u, d_v)}{\min(d_u, d_v)} + \frac{\min(d_u, d_v)}{\max(d_u, d_v)} = \sum_{uv \in E(G)} \frac{d_u}{d_v} + \frac{d_v}{d_u} = \sum_{uv \in E(G)} \frac{d_u^2 + d_v^2}{d_u d_v}$$

where d_v is the degree of a vertex v in G .

The Composition (also called lexicographic product [7]) $G = G_1[G_2]$ of graph G_1 and G_2 with disjoint vertex sets $V(G_1)$ and $V(G_2)$ and edge sets $E(G_1)$ and $E(G_2)$ is the graph with vertex set $V(G_1) \times V(G_2)$ and (u_i, v_j) is adjacent with (u_k, v_l) whenever u_i is adjacent with u_k , or $u_i = u_k$ and v_j is adjacent with v_l .

In [2], the Cartesian product $G_1 \times G_2$ of graph G_1 and G_2 has the vertex set $V(G_1 \times G_2) = V(G_1) \times V(G_2)$ and $(u_i, v_j)(u_k, v_l)$ is an edge of $G_1 \times G_2$ if $u_i = u_k$ and $v_j v_l \in E(G_2)$, or $u_i u_k \in E(G_1)$ and $v_j = v_l$.

For given graph G_1 and G_2 we define their Corona product $G_1 \circ G_2$ as the graph obtained by taking $|V(G_1)|$ copies of G_2 and joining each vertex of the i -th copy with vertex $v_i \in V(G_1)$. Obviously, $|V(G_1 \circ G_2)| = |V(G_1)|(1 + |V(G_2)|)$ and $|E(G_1 \circ G_2)| = |E(G_1)| + |V(G_1)|(|V(G_2)| + |E(G_2)|)$ [4].

A sum $G_1 + G_2$ of two graph G_1 and G_2 with disjoint vertex sets $V(G_1)$ and $V(G_2)$ is the graph on the vertex set $V(G_1) \cup V(G_2)$ and the edge set $E(G_1) \cup E(G_2) \cup \{uv \mid u \in V(G_1), v \in V(G_2)\}$. Hence, the sum of two graph is obtained by connecting each vertex of one graph to each vertex of the other graph, while keeping all edges of both graph.[18].

The Symmetric difference [8] $G_1 \oplus G_2$ of two graph G_1 and G_2 is the graph with vertex set $V(G_1) \times V(G_2)$ and

$$E(G_1 \oplus G_2) = \{(u_1, u_2)(v_1, v_2) \mid u_1 v_1 \in E(G_1) \text{ or } u_2 v_2 \in E(G_2) \text{ but not both}\}.$$

Obviously,

$$|E(G_1 \oplus G_2)| = |E(G_1)||V(G_2)|^2 + |E(G_2)||V(G_1)|^2 - 4|E(G_1)||E(G_2)|$$

$$d_{G_1 \oplus G_2}(u, v) = |V(G_2)|d_{G_1}(u) + |V(G_1)|d_{G_2}(v) - 2d_{G_1}(u)d_{G_2}(v).$$

Some graph operations on Harmonic index are found in [13]. Motivated from this, in this paper, we concentrate on graph operations like join, corona product, cartesian product, composition and symmetric difference of graph are established.

2 Main Results

In this section, we established the graph operations for SDD index.

Theorem 2.1. *Let G_1 and G_2 be two connected graph with order n_1, n_2 , size m_1, m_2 , maximum degree Δ_1, Δ_2 and minimum degree δ_1, δ_2 respectively. Then*

$$SDD(G_1[G_2]) \leq \frac{n_1}{n_2 \delta_1 + \delta_2} (2m_2 n_2 \Delta_1 + M_1(G_2))$$

$$+ \frac{n_2^2}{n_2 \delta_1 + \delta_2} (2m_1 \Delta_2 + n_2 M_1(G_1))$$

Equality hold only if graph is regular.

Proof. Let $V(G_1) = \{u_1, u_2, \dots, u_{n_1}\}$ and $V(G_2) = \{v_1, v_2, \dots, v_{n_2}\}$ be a set of vertex for G_1 and G_2 respectively. By the Definition of the composition of two graph one can see that,

$$|E(G_1[G_2])| = |E(G_1)||V(G_2)|^2 + |E(G_2)||V(G_1)|$$

$$d_{G_1[G_2]}(u, v) = |V(G_2)|d_{G_1}(u) + d_{G_2}(v)$$

Consider,

$$\begin{aligned}
 SDD(G_1[G_2]) &= \sum_{(u_i, v_j), (u_k, v_l) \in E(G_1[G_2]), (u_i, v_j) \neq (u_k, v_l)} \frac{d_{G_1[G_2]}(u_i, v_j)}{d_{G_1[G_2]}(u_k, v_l)} + \frac{d_{G_1[G_2]}(u_k, v_l)}{d_{G_1[G_2]}(u_i, v_j)} \\
 &= \sum_{(u_i, v_j), (u_i, v_l) \in E(G_1[G_2]), j \neq l} \frac{d_{G_1[G_2]}(u_i, v_j)}{d_{G_1[G_2]}(u_i, v_l)} + \frac{d_{G_1[G_2]}(u_i, v_l)}{d_{G_1[G_2]}(u_i, v_j)} \\
 &\quad + \sum_{(u_i, v_j), (u_k, v_l) \in E(G_1[G_2]), i \neq k} \frac{d_{G_1[G_2]}(u_i, v_j)}{d_{G_1[G_2]}(u_k, v_l)} + \frac{d_{G_1[G_2]}(u_k, v_l)}{d_{G_1[G_2]}(u_i, v_j)} \\
 &= \sum_{u_i \in V(G_1)} \sum_{v_j, v_l \in E(G_2)} \frac{|V(G_2)|d_{G_1}(u_i) + d_{G_2}(v_j)}{|V(G_2)|d_{G_1}(u_i) + d_{G_2}(v_l)} + \frac{|V(G_2)|d_{G_1}(u_i) + d_{G_2}(v_l)}{|V(G_2)|d_{G_1}(u_i) + d_{G_2}(v_j)} \\
 &\quad + \sum_{u_i, u_k \in E(G_1)} \sum_{v_j \in V(G_2)} \sum_{v_l \in V(G_2)} \frac{|V(G_2)|d_{G_1}(u_i) + d_{G_2}(v_j)}{|V(G_2)|d_{G_1}(u_k) + d_{G_2}(v_l)} + \frac{|V(G_2)|d_{G_1}(u_k) + d_{G_2}(v_l)}{|V(G_2)|d_{G_1}(u_i) + d_{G_2}(v_j)} \\
 &= n_1 \sum_{v_j, v_l \in E(G_2)} \frac{n_2 d_{G_1}(u_i) + d_{G_2}(v_j)}{n_2 d_{G_1}(u_i) + d_{G_2}(v_l)} + \frac{n_2 d_{G_1}(u_i) + d_{G_2}(v_l)}{n_2 d_{G_1}(u_i) + d_{G_2}(v_j)} \\
 &\quad + n_2^2 \sum_{u_i, u_k \in E(G_1)} \frac{n_2 d_{G_1}(u_i) + d_{G_2}(v_j)}{n_2 d_{G_1}(u_k) + d_{G_2}(v_l)} + \frac{n_2 d_{G_1}(u_k) + d_{G_2}(v_l)}{n_2 d_{G_1}(u_i) + d_{G_2}(v_j)} \\
 &\leq n_1 \sum_{v_j, v_l \in E(G_2)} \frac{n_2 \Delta_1 + d_{G_2}(v_j)}{n_2 \delta_1 + \delta_2} + \frac{n_2 \Delta_1 + d_{G_2}(v_l)}{n_2 \delta_1 + \delta_2} \\
 &\quad + n_2^2 \sum_{u_i, u_k \in E(G_1)} \frac{n_2 d_{G_1}(u_i) + \Delta_2}{n_2 \delta_1 + \delta_2} + \frac{n_2 d_{G_1}(u_k) + \Delta_2}{n_2 \delta_1 + \delta_2} \\
 &= \frac{n_1}{n_2 \delta_1 + \delta_2} \sum_{v_j, v_l \in E(G_2)} (2n_2 \Delta_1 + d_{G_2}(v_j) + d_{G_2}(v_l)) \\
 &\quad + \frac{n_2^2}{n_2 \delta_1 + \delta_2} \sum_{u_i, u_k \in E(G_1)} (2\Delta_2 + n_2(d_{G_1}(u_i) + d_{G_1}(u_k))) \\
 &= \frac{n_1}{n_2 \delta_1 + \delta_2} (2m_2 n_2 \Delta_1 + M_1(G_2)) + \frac{n_2^2}{n_2 \delta_1 + \delta_2} (2m_1 \Delta_2 + n_2 M_1(G_1)).
 \end{aligned}$$

□

Corollary 2.2. Let G_i for $i \in \{1, 2\}$, be a cayley graph of nilpotent matrix group of length one. Then

$$SDD(G_1[G_2]) = 4n^2(n + 1) \text{ for } n \geq 7.$$

Proof. Result is directly from the theorem 2.1. Since For $n \geq 7$, cayley graph of nilpotent matrix group of length one with n vertices has $2n$ edges and degree of each vertices is 4. Also $M_1(G_i) = 16n$. □

Theorem 2.3. Let G_1 and G_2 be two connected graph with order n_1, n_2 and size m_1, m_2 respectively. Then

$$SDD(G_1 \times G_2) \leq \frac{2(\Delta_1 + \Delta_2)}{\delta_1 + \delta_2} (n_1 m_2 + n_2 m_1).$$

Equality hold only if graph is regular.

Proof. Let $V(G_1) = \{u_1, u_2, \dots, u_{n_1}\}$ and $V(G_2) = \{v_1, v_2, \dots, v_{n_2}\}$ be a set of vertex for G_1 and G_2 respectively. By the Definition of the cartesian product of two graph one can see that,

$$\begin{aligned}
 |E(G_1 \times G_2)| &= |E(G_1)||V(G_2)| + |E(G_2)||V(G_1)| \\
 d_{G_1 \times G_2}(u, v) &= d_{G_1}(u) + d_{G_2}(v).
 \end{aligned}$$

Consider,

$$\begin{aligned}
 SDD(G_1 \times G_2) &= \sum_{(u_i, v_j), (u_k, v_l) \in E(G_1 \times G_2), (u_i, v_j) \neq (u_k, v_l)} \left[\frac{d_{G_1 \times G_2}(u_i, v_j)}{d_{G_1 \times G_2}(u_k, v_l)} + \frac{d_{G_1 \times G_2}(u_k, v_l)}{d_{G_1 \times G_2}(u_i, v_j)} \right] \\
 &= \sum_{(u_i, v_j), (u_i, v_l) \in E(G_1 \times G_2), v_j, v_l \in E(G_2)} \left[\frac{d_{G_1 \times G_2}(u_i, v_j)}{d_{G_1 \times G_2}(u_i, v_l)} + \frac{d_{G_1 \times G_2}(u_i, v_l)}{d_{G_1 \times G_2}(u_i, v_j)} \right] \\
 &+ \sum_{(u_i, v_j), (u_k, v_j) \in E(G_1 \times G_2), u_i, u_k \in E(G_1)} \left[\frac{d_{G_1 \times G_2}(u_i, v_j)}{d_{G_1 \times G_2}(u_k, v_j)} + \frac{d_{G_1 \times G_2}(u_k, v_j)}{d_{G_1 \times G_2}(u_i, v_j)} \right] \\
 &= \sum_{u_i \in V(G_1)} \sum_{v_j, v_l \in E(G_2)} \left[\frac{d_{G_1}(u_i) + d_{G_2}(v_j)}{d_{G_1}(u_i) + d_{G_2}(v_l)} + \frac{d_{G_1}(u_i) + d_{G_2}(v_l)}{d_{G_1}(u_i) + d_{G_2}(v_j)} \right] \\
 &+ \sum_{v_j \in V(G_2)} \sum_{u_i, u_k \in E(G_1)} \left[\frac{d_{G_1}(u_i) + d_{G_2}(v_j)}{d_{G_1}(u_k) + d_{G_2}(v_j)} + \frac{d_{G_1}(u_k) + d_{G_2}(v_j)}{d_{G_1}(u_i) + d_{G_2}(v_j)} \right] \\
 &\leq n_1 \sum_{v_j, v_l \in E(G_2)} \left[\frac{\Delta_1 + \Delta_2}{\delta_1 + \delta_2} + \frac{\Delta_1 + \Delta_2}{\delta_1 + \delta_2} \right] \\
 &+ n_2 \sum_{u_i, u_k \in E(G_1)} \left[\frac{\Delta_1 + \Delta_2}{\delta_1 + \delta_2} + \frac{\Delta_1 + \Delta_2}{\delta_1 + \delta_2} \right] \\
 &= 2n_1m_2 \left[\frac{\Delta_1 + \Delta_2}{\delta_1 + \delta_2} \right] + 2n_2m_1 \left[\frac{\Delta_1 + \Delta_2}{\delta_1 + \delta_2} \right] \\
 &= 2 \left[\frac{\Delta_1 + \Delta_2}{\delta_1 + \delta_2} \right] (n_1m_2 + n_2m_1).
 \end{aligned}$$

Equality hold only if graph is regular. □

Corollary 2.4. Let G_i for $i \in \{1, 2\}$, be a cayley graph of nilpotent matrix group of length one. Then

$$SDD(G_1 \times G_2) = 8n^2 \text{ for } n \geq 7.$$

Theorem 2.5. For $i \in \{1, 2\}$, let G_i be a graph of minimum degree δ_i , maximum degree Δ_i , order n_i and size m_i . Then

$$SDD(G_1 \circ G_2) \leq 2m_1 \left(\frac{\Delta_1 + n_2}{\delta_1 + n_2} \right) + 2m_2n_1 \left(\frac{\Delta_2 + 1}{\delta_2 + 1} \right) + n_1n_2 \left(\frac{\Delta_1 + n_2}{\delta_2 + 1} + \frac{\Delta_2 + 1}{\delta_1 + n_2} \right).$$

Equality hold only if graph is regular.

Proof. The edges of $G_1 \circ G_2$ are partitioned into three subsets E_1, E_2 and E_3 as follows

$$\begin{aligned}
 E_1 &= \{e \in E(G_1 \circ G_2), e \in E(G_1)\} \\
 E_2 &= \{e \in E(G_1 \circ G_2), e \in E(G_{2i}), i = 1, 2, \dots, |V(G_1)|\} \\
 E_3 &= \{e \in E(G_1 \circ G_2), e = uv, u \in V(G_{2i}), i = 1, 2, \dots, |V(G_1)| \text{ and } v \in V(G_1)\}.
 \end{aligned}$$

and if u is a vertex of $G_1 \circ G_2$, then

$$d_{G_1 \circ G_2}(u) = \begin{cases} d_{G_1}(u) + |V(G_2)| & \text{if } u \in V(G_1) \\ d_{G_2}(u) + 1 & \text{if } u \in V(G_2) \end{cases}$$

Let $G_1 = (V_i, E_i), i \in \{1, 2\}$ and let $G_1 \circ G_2 = (V, E)$ we have

$$\begin{aligned}
 SDD(G_1 \circ G_2) &= \sum_{uv \in E(G_1 \circ G_2)} \left(\frac{d_{G_1 \circ G_2}(u)}{d_{G_1 \circ G_2}(v)} + \frac{d_{G_1 \circ G_2}(v)}{d_{G_1 \circ G_2}(u)} \right) \\
 &= Q_1 + Q_2 + Q_3.
 \end{aligned}$$

where

$$\begin{aligned}
 Q_1 &= \sum_{uv \in E_1} \left(\frac{d_{G_1}(u) + n_2}{d_{G_1}(v) + n_2} + \frac{d_{G_1}(v) + n_2}{d_{G_1}(u) + n_2} \right) \\
 &\leq \sum_{uv \in E_1} \left(\frac{\Delta_1 + n_2}{\delta_1 + n_2} + \frac{\Delta_1 + n_2}{\delta_1 + n_2} \right) = 2m_1 \left(\frac{\Delta_1 + n_2}{\delta_1 + n_2} \right) \\
 Q_2 &= n_1 \sum_{uv \in E_2} \left(\frac{d_{G_2}(u) + 1}{d_{G_1}(v) + 1} + \frac{d_{G_1}(v) + 1}{d_{G_1}(u) + 1} \right) \\
 &\leq n_1 m_2 \left(\frac{\Delta_2 + 1}{\delta_2 + 1} + \frac{\Delta_2 + 1}{\delta_2 + 1} \right) = 2n_1 m_2 \left(\frac{\Delta_2 + 1}{\delta_2 + 1} \right) \\
 Q_3 &= \sum_{uv \in E_3, u \in V_1 \text{ and } v \in V_2} \left(\frac{d_{G_1}(u) + n_2}{d_{G_2}(v) + 1} + \frac{d_{G_2}(v) + 1}{d_{G_1}(u) + n_2} \right) \\
 &\leq n_1 n_2 \left(\frac{\Delta_1 + n_2}{\delta_2 + 1} + \frac{\Delta_2 + 1}{\delta_1 + n_2} \right).
 \end{aligned}$$

Using Q_1 to Q_3 in $SDD(G_1 \circ G_2)$, we get

$$SDD(G_1 \circ G_2) \leq 2m_1 \left(\frac{\Delta_1 + n_2}{\delta_1 + n_2} \right) + 2m_2 n_1 \left(\frac{\Delta_2 + 1}{\delta_2 + 1} \right) + n_1 n_2 \left(\frac{\Delta_1 + n_2}{\delta_2 + 1} + \frac{\Delta_2 + 1}{\delta_1 + n_2} \right).$$

□

Corollary 2.6. Let G_i for $i \in \{1, 2\}$, be a cayley graph of nilpotent matrix group of length one. Then

$$SDD(G_1 \circ G_2) = 4n^2 + 4n + n^2 \left(\frac{n^2 + 8n + 41}{5n + 20} \right) \text{ for } n \geq 7.$$

Theorem 2.7. Let G_1 and G_2 be two connected graph with order n_1, n_2 and size m_1, m_2 respectively. Then

$$SDD(G_1 + G_2) \leq 2m_1 \left(\frac{\Delta_1 + n_2}{\delta_1 + n_2} \right) + 2m_2 \left(\frac{\Delta_2 + n_1}{\delta_2 + n_1} \right) + n_1 n_2 \left(\frac{\Delta_1 + n_2}{\delta_2 + n_1} + \frac{\Delta_2 + n_1}{\delta_1 + n_2} \right).$$

Proof. Let $V(G_1) = \{u_1, u_2, \dots, u_{n_1}\}$ and $V(G_2) = \{v_1, v_2, \dots, v_{n_2}\}$ be a set of vertex for G_1 and G_2 respectively. By the Definition of the join of two graph one can see that, if u is a vertex of $G_1 + G_2$, then

$$d_{G_1+G_2}(u) = \begin{cases} d_{G_1}(u) + |V(G_2)| & \text{if } u \in V(G_1) \\ d_{G_2}(u) + |V(G_1)| & \text{if } u \in V(G_2) \end{cases}$$

Therefore,

$$\begin{aligned}
 SDD(G_1 + G_2) &= \sum_{uv \in E(G_1+G_2)} \left(\frac{d_{G_1+G_2}(u)}{d_{G_1+G_2}(v)} + \frac{d_{G_1+G_2}(v)}{d_{G_1+G_2}(u)} \right) \\
 &= \sum_{uv \in E(G_1)} \left(\frac{d_{G_1}(u) + n_2}{d_{G_1}(v) + n_2} + \frac{d_{G_1}(v) + n_2}{d_{G_1}(u) + n_2} \right) + \sum_{uv \in E(G_2)} \left(\frac{d_{G_2}(u) + n_1}{d_{G_2}(v) + n_1} + \frac{d_{G_2}(v) + n_1}{d_{G_2}(u) + n_1} \right) \\
 &\quad + \sum_{u \in V(G_1), v \in V(G_2)} \left(\frac{d_{G_1}(u) + n_2}{d_{G_2}(v) + n_1} + \frac{d_{G_2}(v) + n_1}{d_{G_1}(u) + n_2} \right) \\
 &\leq 2m_1 \left(\frac{\Delta_1 + n_2}{\delta_1 + n_2} \right) + 2m_2 \left(\frac{\Delta_2 + n_1}{\delta_2 + n_1} \right) + n_1 n_2 \left(\frac{\Delta_1 + n_2}{\delta_2 + n_1} + \frac{\Delta_2 + n_1}{\delta_1 + n_2} \right).
 \end{aligned}$$

□

Corollary 2.8. Let G_i for $i \in \{1, 2\}$, be a cayley graph of nilpotent matrix group of length one. Then

$$SDD(G_1 + G_2) = 2n^2 + 8n \text{ for } n \geq 7.$$

Theorem 2.9. For $i \in \{1, 2\}$, let G_i be a graph of maximum degree Δ_i , minimum degree δ_i order n_i and size m_i . Then

$$SDD(G_1 \oplus G_2) \leq 2(n_2^2m_1 + n_1^2m_2 - 4m_1m_2) \frac{n_2\Delta_1 + n_1\Delta_2 - 2\delta_1\delta_2}{n_2\delta_1 + n_1\delta_2 - 2\Delta_1\Delta_2}.$$

Proof. Let $V(G_1) = \{u_1, u_2, \dots, u_{n_1}\}$ and $V(G_2) = \{v_1, v_2, \dots, v_{n_2}\}$ be a set of vertex for G_1 and G_2 respectively.

Consider,

$$\begin{aligned} SDD(G_1 \oplus G_2) &= \sum_{(u_i, v_j), (u_k, v_l) \in E(G_1 \oplus G_2)} \frac{d_{G_1 \oplus G_2}(u_i, v_j)}{d_{G_1 \oplus G_2}(u_k, v_l)} + \frac{d_{G_1 \oplus G_2}(u_k, v_l)}{d_{G_1 \oplus G_2}(u_i, v_j)} \\ &= \sum_{v_j \in V(G_2)} \sum_{v_l \in V(G_2)} \sum_{u_i, u_k \in E(G_1)} \frac{d_{G_1 \oplus G_2}(u_i, v_j)}{d_{G_1 \oplus G_2}(u_k, v_l)} + \frac{d_{G_1 \oplus G_2}(u_k, v_l)}{d_{G_1 \oplus G_2}(u_i, v_j)} \\ &\quad + \sum_{u_i \in V(G_1)} \sum_{u_k \in V(G_1)} \sum_{v_j, v_l \in E(G_2)} \frac{d_{G_1 \oplus G_2}(u_i, v_j)}{d_{G_1 \oplus G_2}(u_k, v_l)} + \frac{d_{G_1 \oplus G_2}(u_k, v_l)}{d_{G_1 \oplus G_2}(u_i, v_j)} \\ &\quad - \sum_{u_i, u_k \in E(G_1)} \sum_{v_j, v_l \in E(G_1)} \frac{d_{G_1 \oplus G_2}(u_i, v_j)}{d_{G_1 \oplus G_2}(u_k, v_l)} + \frac{d_{G_1 \oplus G_2}(u_k, v_l)}{d_{G_1 \oplus G_2}(u_i, v_j)} \dots\dots(1) \end{aligned}$$

As per the definition of the Symmetric difference of a graph

$$\begin{aligned} \frac{d_{G_1 \oplus G_2}(u_i, v_j)}{d_{G_1 \oplus G_2}(u_k, v_l)} + \frac{d_{G_1 \oplus G_2}(u_k, v_l)}{d_{G_1 \oplus G_2}(u_i, v_j)} &= \frac{n_2d_{G_1}(u_i) + n_1d_{G_2}(v_j) - 2d_{G_1}(u_i)d_{G_2}(v_j)}{n_2d_{G_1}(u_k) + n_1d_{G_2}(v_l) - 2d_{G_1}(u_k)d_{G_2}(v_l)} \\ &\quad + \frac{n_2d_{G_1}(u_k) + n_1d_{G_2}(v_l) - 2d_{G_1}(u_k)d_{G_2}(v_l)}{n_2d_{G_1}(u_i) + n_1d_{G_2}(v_j) - 2d_{G_1}(u_i)d_{G_2}(v_j)} \\ &\leq \frac{n_2\Delta_1 + n_1\Delta_2 - 2\delta_1\delta_2}{n_2\delta_1 + n_1\delta_2 - 2\Delta_1\Delta_2} + \frac{n_2\Delta_1 + n_1\Delta_2 - 2\delta_1\delta_2}{n_2\delta_1 + n_1\delta_2 - 2\Delta_1\Delta_2} \\ &= 2 \frac{n_2\Delta_1 + n_1\Delta_2 - 2\delta_1\delta_2}{n_2\delta_1 + n_1\delta_2 - 2\Delta_1\Delta_2} \dots\dots(2) \end{aligned}$$

From equation (1) and (2) we get

$$\begin{aligned} SDD(G_1 \oplus G_2) &\leq n_2^2m_1 \frac{2(n_2\Delta_1 + n_1\Delta_2 - 2\delta_1\delta_2)}{n_2\delta_1 + n_1\delta_2 - 2\Delta_1\Delta_2} \\ &\quad + n_1^2m_2 \frac{2(n_2\Delta_1 + n_1\Delta_2 - 2\delta_1\delta_2)}{n_2\delta_1 + n_1\delta_2 - 2\Delta_1\Delta_2} \\ &\quad - 4m_1m_2 \frac{2(n_2\Delta_1 + n_1\Delta_2 - 2\delta_1\delta_2)}{n_2\delta_1 + n_1\delta_2 - 2\Delta_1\Delta_2} \\ &= 2(n_2^2m_1 + n_1^2m_2 - 4m_1m_2) \frac{n_2\Delta_1 + n_1\Delta_2 - 2\delta_1\delta_2}{n_2\delta_1 + n_1\delta_2 - 2\Delta_1\Delta_2}. \end{aligned}$$

□

Corollary 2.10. Let G_i for $i \in \{1, 2\}$, be a cayley graph of nilpotent matrix group of length one. Then

$$SDD(G_1 + G_2) = 8n^2(n - 4) \text{ for } n \geq 7.$$

References

[1] V. Alexander, Upper and lower bounds of symmetric division deg index, *Iranian Journal of Mathematical Chemistry*, 5,2, (2014) 91–98.

- [2] Das. K. C , Xu. K ,Cangul. I.N, Cevik. A.S and Graovac.A, On the Harary index of graph operations, *Journal of Inequalities and Applications*,1(2013), 1-16.
- [3] J. Devillers and A. T. Balaban (Eds.), **Topological Indices and Related Descriptors in QSAR and QSPR**, Gordon and Breach, Amsterdam, (1999).
- [4] G.H. Fath-Tabar, A. Hamzeh and S. Hossein-Zadeh, GA2 index of some graph operations, *Filomat*, **24**:1(2010), 210–218.
- [5] C. K. Gupta, V. Lokesha, Shwetha B. Shetty, On the graph of nilpotent matrix group of length one, *Discrete Mathematics, Algorithms and applications* **8(1)**(2016), 1-31. DOI: 10.1142/S1793830916500099.
- [6] C. K. Gupta, V. Lokesha, Shwetha B. Shetty, On the Symmetric division deg index of graph, *South East Asian Journal Of Mathematics* **41(1)** (2016), 1-23.
- [7] F. Harary, **Graph theory**, Reading, MA: Addison-Wesley, (1994).
- [8] M.H. Khalifeha, H. Yousefi-Azari and A. R. Ashrafi, The Hyper-Wiener index of graph operations, *Computers and Mathematics with Applications*, **56:5** (2008), 1402–1407.
- [9] M.H. Khalifeha, H. Yousefi-Azari and A. R. Ashrafi, The first and second Zagreb indices of some graph operations, *Discrete Applied Mathematics*, **157:4**(2009), 804–811.
- [10] V. Lokesha, Shwetha B. S, Ranjini P.S, Cangul. Ismail N and Cevik Ahmet S, New bounds for Randic and GA indices, *Journal of Inequalities and Applications*, **180:1**, (2013), 1-7.
- [11] Shwetha B. Shetty, V. Lokesha, P. S. Ranjini and K. C. Das, Computing some topological indices of Smart polymer, *Digest Journal of Nanomaterials and Biostructures*, **7:3**(2012), 1097–1102.
- [12] B. S. Shwetha, V. Lokesha, A. Bayad and P. S. Ranjini, A Comparative Study of Topological Indices and Molecular Weight of Some Carbohydrates, *Journal of the Indian Academy of Mathematics*, **34(2)** (2012), 627-636.
- [13] B. S. Shwetha, V. Lokesha and P. S. Ranjini, On The Harmonic Index of Graph Operations, *Transactions on Combinatorics*, **4(4)**(2015), 5-14 .
- [14] D. Suprunenko, *Mairix groups*, Transl. Math. Monogr., vol. 45, **American Mathematical Society**, Providence, RI, (1976).
- [15] Shirdel. G. H., H. Rezapour, and A. M. Sayadi, The Hyper-Zagreb index of Graph Operations, *Iranian Journal of Mathematical Chemistry* **4.2** (2013), 213-220.
- [16] Vukicevic. D and M. Gažperov, Bond additive modeling 1. Adriatic indices, *Croatica chemica acta* **83.3** (2010), 243-260.
- [17] Vukicevic. D, Bond Additive Modeling 2. Mathematical properties of Max-min rodeg index, *Croatica chemica acta* **83.3**(2010), 261-273.
- [18] Yero. Ismael. G and Rodríguez-Velázquez. Juan A, *On the Randić index of Corona Product graph*, International Scholarly Research Notices, (2011).

Author information

C. K. Gupta¹, V. Lokesha^{2*}, Shwetha B. S³ and Ranjini P. S⁴,

¹Department of Mathematics, University of Manitoba, Winnipeg, R3T 2N2, Canada.

²Department of Studies in Mathematics, V. S. K University, Ballari-583105, India.

^{3,4}Department of Mathematics, Don Bosco Institute of Technology, Bangalore-78, India.,.

E-mail: Chander.Gupta@umanitoba.ca; v.lokesha@gmail.com; shwethu17@gmail.com;
ranjini_p_s@yahoo.com

Received: June 12, 2015.

Accepted: March 21, 2016.