

Study on double integral operator associated with generalized Bessel-Maitland function

Musharraf Ali, Waseem A. Khan and Idrees A. Khan

Communicated by Jose Luis Lopez-Bonilla

MSC 2010 Classifications: 33C45, 33C60, 33E12.

Keywords and phrases: Generalized Bessel-Maitland function, Generalized (Wright) hypergeometric function and integrals, Mittag-Leffler function.

Abstract In this article, we establish double integral operator involving generalized Bessel-Maitland function defined by Khan et al. [9], which is expressed in terms of generalized (Wright) hypergeometric function. Some special cases are deduced.

1 Introduction

In last decade many authors namely, Choi and Agarwal [1, 2, 3], Khan et al. [7-10], Ghayasuddin and Khan [4] have introduced integral formulae associated with Bessel function. It has a wide application in the problem of Physics, Chemistry, Biology, Engineering and Applied Sciences. The theory of Bessel functions is closely associated with the theory of certain types of differential equations (see [21]).

The Bessel-Maitland function $J_\nu^\mu(z)$ [13;Eq.(8.3)] is defined by means of the following series representation:

$$J_\nu^\mu(z) = \sum_{n=0}^{\infty} \frac{(-z)^n}{n! \Gamma(\mu n + \nu + 1)} = \phi(\mu, \nu + 1; -z). \quad (1.1)$$

Singh et al. [20] introduced the following generalization of Bessel-Maitland function:

$$J_{\nu,q}^{\mu,\gamma} = \sum_{n=0}^{\infty} \frac{(\gamma)_{qn} (-z)^n}{\Gamma(\mu n + \nu + 1) n!}, \quad (1.2)$$

where $\mu, \nu, \gamma \in \mathbb{C}$, $\operatorname{Re}(\mu) \geq 0$, $\operatorname{Re}(\nu) \geq -1$, $\operatorname{Re}(\gamma) \geq 0$ and $q \in (0, 1) \cup \mathbb{N}$ and $(\gamma)_0 = 1$, $(\gamma)_{qn} = \frac{\Gamma(\gamma+qn)}{\Gamma(\gamma)}$ denotes the generalized Pochhammer symbol.

Recently, Ghayasuddin and Khan [4] introduced and investigated generalized Bessel-Maitland function defined as

$$J_{\nu,\gamma,\delta}^{\mu,q,p}(z) = \sum_{n=0}^{\infty} \frac{(\gamma)_{qn} (-z)^n}{\Gamma(\mu n + \nu + 1) (\delta)_{pn}}, \quad (1.3)$$

where $\mu, \nu, \gamma, \delta \in \mathbb{C}$, $\operatorname{Re}(\mu) \geq 0$, $\operatorname{Re}(\nu) \geq -1$, $\operatorname{Re}(\gamma) \geq 0$, $\operatorname{Re}(\delta) \geq 0$; $p, q > 0$ and $q < \operatorname{Re}(\alpha) + p$.

In particular, Khan et al. [10] introduced and investigated a new extension of Bessel-Maitland function as follows:

$$J_{\alpha,\beta,\nu,\sigma,\delta,p}^{\mu,\rho,\gamma,q}(z) = \sum_{n=0}^{\infty} \frac{(\mu)_{pn} (\gamma)_{qn} (-z)^n}{\Gamma(n\beta + \alpha + 1) (\delta)_{pn} (\nu)_{n\sigma}}, \quad (1.4)$$

where $\alpha, \beta, \mu, \rho, \nu, \gamma, \delta, \sigma \in \mathbb{C}$; $\operatorname{Re}(\sigma) \geq 0$, $\operatorname{Re}(\beta) \geq 0$, $\operatorname{Re}(\rho) \geq 0$, $\operatorname{Re}(\mu) \geq 0$, $\operatorname{Re}(\nu) \geq 0$, $\operatorname{Re}(\alpha) \geq -1$, $\operatorname{Re}(\gamma) \geq 0$, $\operatorname{Re}(\delta) \geq 0$; $p, q > 0$, and $q < \operatorname{Re}(\alpha) + p$.

The Mittag-Leffler (Swedish Mathematician) [12] introduced the function $E_\alpha(z)$ and defined it as:

$$E_\alpha(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + 1)}, \quad (1.5)$$

where $z \in \mathbb{C}$ and $\Gamma(s)$ is the Gamma function; $\alpha \geq 0$.

The Mittag-Leffler function is a direct generalization of $\exp(z)$ in which $\alpha = 1$. Mittag-Leffler function naturally occurs as the solution of fractional order differential equation or fractional order integral equations.

A generalization of $E_\alpha(z)$ was studied by Wiman [22] where he defined the function $E_{\alpha,\beta}(z)$ as:

$$E_{\alpha,\beta}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + \beta)}, \quad (1.6)$$

where $\alpha, \beta \in \mathbb{C}; \Re(\alpha) > 0, \Re(\beta) > 0$ which is also known as Mittag-Leffler function or Wiman's function.

In (1903), Prabhakar [14] introduced the function $E_{\alpha,\beta}^\gamma(z)$ as:

$$E_{\alpha,\beta}^\gamma(z) = \sum_{n=0}^{\infty} \frac{(\gamma)_n}{\Gamma(\alpha n + \beta)} \frac{z^n}{n!}, \quad (1.7)$$

where $\alpha, \beta, \gamma \in \mathbb{C}; \Re(\alpha) > 0, \Re(\beta) > 0, \Re(\gamma) > 0$.

Shukla and Prajapati [19] defined and investigated the function $E_{\alpha,\beta}^{\gamma,q}(z)$ as:

$$E_{\alpha,\beta}^{\gamma,q}(z) = \sum_{n=0}^{\infty} \frac{(\gamma)_{qn}}{\Gamma(\alpha n + \beta)} \frac{z^n}{n!}, \quad (1.8)$$

where $\alpha, \beta, \gamma \in \mathbb{C}; \Re(\alpha) > 0, \Re(\beta) > 0, \Re(\gamma) > 0, q \in (0, 1) \cup \mathbb{N}$ and $(\gamma)_{qn} = \frac{\Gamma(\gamma + qn)}{\Gamma(\gamma)}$ denotes the generalized Pochhammer symbol which in particular reduces to $q^{qn} \prod_{r=1}^q \left(\frac{\gamma + r - 1}{q} \right)_n$ if $q \in \mathbb{N}$.

Salim [18] introduced a new generalized Mittag-Leffler function and defined it as:

$$E_{\alpha,\beta}^{\gamma,\delta}(z) = \sum_{n=0}^{\infty} \frac{(\gamma)_n}{\Gamma(\alpha n + \beta)} \frac{z^n}{(\delta)_n}, \quad (1.9)$$

where $\alpha, \beta, \gamma, \delta \in \mathbb{C}; \Re(\alpha) > 0, \Re(\beta) > 0, \Re(\gamma) > 0, \Re(\delta) > 0$.

Salim and Faraj [17] introduced the generalized Mittag-Leffler function $E_{\alpha,\beta,p}^{\gamma,\delta,q}(z)$ which is defined as:

$$E_{\alpha,\beta,p}^{\gamma,\delta,q}(z) = \sum_{n=0}^{\infty} \frac{(\gamma)_{qn}}{\Gamma(\alpha n + \beta)} \frac{z^n}{(\delta)_{pn}}, \quad (1.10)$$

where $\alpha, \beta, \gamma, \delta \in \mathbb{C}, \min\{\Re(\alpha) > 0, \Re(\beta) > 0, \Re(\gamma) > 0, \Re(\delta) > 0\} > 0; p, q > 0$ and $q < \Re\alpha + p$.

In (2016), Khan and Ahmed [11] introduced and investigated generalized Mittag Leffler function is defined as

$$E_{\alpha,\beta,\nu,\sigma,\delta,p}^{\mu,\rho,\gamma,q}(z) = \sum_{n=0}^{\infty} \frac{(\mu)_{pn} (\gamma)_{qn}}{\Gamma(\alpha n + \beta) (\nu)_{\sigma n} (\delta)_{pn}} z^n, \quad (1.11)$$

where $\alpha, \beta, \gamma, \delta, \mu, \nu, \rho, \sigma \in \mathbb{C}$, $\Re(\alpha) > 0, \Re(\beta) > 0, \Re(\gamma) > 0, \Re(\delta) > 0, \Re(\mu) > 0, \Re(\nu) > 0, \Re(\rho) > 0, \Re(\sigma) > 0, p, q > 0$ and $q \leq \Re(\alpha) + p$.

The generalization of the generalized hypergeometric series ${}_pF_q$ is due to Fox [6] and Wright (see [23, 24]) who studied the asymptotic expansion of the generalized (Wright) hypergeometric function defined by (see [16, p.21]; see also [15]):

$${}_p\Psi_q \left[\begin{array}{l} (\alpha_1, A_1), \dots, (\alpha_p, A_p); \\ (\beta_1, B_1), \dots, (\beta_q, B_q); \end{array} z \right] = \sum_{k=0}^{\infty} \frac{\prod_{j=1}^p \Gamma(\alpha_j + A_j k)}{\prod_{j=1}^q \Gamma(\beta_j + B_j k)} \frac{z^k}{k!}, \quad (1.12)$$

where the coefficients A_1, \dots, A_p and B_1, \dots, B_q are positive real numbers such that

$$(i) \quad 1 + \sum_{j=1}^q B_j - \sum_{j=1}^p A_j > 0 \text{ and } 0 < |z| < \infty; \quad z \neq 0. \quad (1.13)$$

$$(ii) \quad 1 + \sum_{j=1}^q B_j - \sum_{j=1}^p A_j = 0 \text{ and } 0 < |z| < A_1^{-A_1} \dots A_p^{-A_p} B_1^{B_1} \dots B_q^{B_q}. \quad (1.14)$$

A special case of (1.12) is

$${}_p\Psi_q \left[\begin{array}{l} (\alpha_1, 1), \dots, (\alpha_p, 1); \\ (\beta_1, 1), \dots, (\beta_q, 1); \end{array} z \right] = \frac{\prod_{j=1}^p \Gamma(\alpha_j)}{\prod_{j=1}^q \Gamma(\beta_j)} {}_pF_q \left[\begin{array}{l} \alpha_1, \dots, \alpha_p; \\ \beta_1, \dots, \beta_q; \end{array} z \right], \quad (1.15)$$

where ${}_pF_q$ is the generalized hypergeometric series defined by [15]:

$$\begin{aligned} {}_pF_q \left[\begin{array}{l} \alpha_1, \dots, \alpha_p; \\ \beta_1, \dots, \beta_q; \end{array} z \right] &= \sum_{n=0}^{\infty} \frac{(\alpha_1)_n \cdots (\alpha_p)_n}{(\beta_1)_n \cdots (\beta_q)_n} \frac{z^n}{n!} \\ &= {}_pF_q(\alpha_1, \dots, \alpha_p; \beta_1, \dots, \beta_q; z), \end{aligned} \quad (1.16)$$

where $(\lambda)_n$ is the Pochhammer's symbol [15].

Furthermore, we also state here the following interesting and useful result stated by Edward [5, p.445]:

$$\int_0^1 \int_0^1 y^\lambda (1-x)^{\lambda-1} (1-y)^{\mu-1} (1-xy)^{1-\lambda-\mu} dx = \frac{\Gamma(\lambda)\Gamma(\mu)}{\Gamma(\lambda+\mu)}, \quad (1.17)$$

provided $0 < \Re(\mu) < \Re(\lambda)$.

2 Unified integral operator involving generalized Bessel-Maitland function

Theorem 2.1. If $\alpha, \beta, \gamma, \mu, \nu, \sigma, \rho, \delta, \lambda \in \mathbb{C}$, $\sigma + \eta + p - \rho - q > 0, \Re(\alpha) > 0, \Re(\beta) > 0, \Re(\gamma) > 0, \Re(\mu) > 0, \Re(\lambda) \geq -1, \Re(\eta) > 0, \Re(\nu) > 0, \Re(\delta) > 0, \Re(\sigma) > 0, \Re(\rho) > 0, ; p, q > 0$ and $q < \Re(\alpha) + p.$, then

$$\begin{aligned} &\int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} J_{\eta, \lambda, \nu, \sigma, \delta, p}^{\mu, \rho, \gamma, q} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy \\ &= \frac{\Gamma(\nu)\Gamma(\delta)}{\Gamma(\mu)\Gamma(\gamma)} {}_5\Psi_4 \left[\begin{array}{l} (\mu, \rho), (\gamma, q), (\alpha, 1), (\beta, 1), (1, 1); \\ (\nu, \sigma), (\delta, p), (\lambda + 1, \eta), (\alpha + \beta, 2); \end{array} -a \right]. \end{aligned} \quad (2.1)$$

Proof. To establish our main result (2.1), we denote the left-hand side of (2.1) by I and then by using (1.4), we have:

$$\begin{aligned} I &= \int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} \\ &\quad \times \sum_{n=0}^{\infty} \frac{(\mu)_{pn} (\gamma)_{qn}}{\Gamma(\eta n + \lambda + 1) (\delta)_{pn} (\nu)_{\sigma n}} \left[\frac{-ay(1-x)(1-y)}{(1-xy)^2} \right]^n dx dy. \end{aligned} \quad (2.2)$$

Now changing the order of integration and summation, (which is guaranteed under the given condition), to get:

$$I = \frac{\Gamma(\delta)\Gamma(\nu)}{\Gamma(\mu)\Gamma(\gamma)} \sum_{n=0}^{\infty} \frac{\Gamma(\mu + \rho n)\Gamma(\gamma + q n)\Gamma(\alpha + n)\Gamma(\beta + n)\Gamma(1+n)}{\Gamma(\nu + \sigma n)\Gamma(\eta n + \lambda + 1)\Gamma(\delta + p n)\Gamma(\alpha + \beta + 2n)} \frac{(-a)^n}{n!}. \quad (2.3)$$

Finally, summing up the above series with the help of (1.12), we easily arrive at the right-hand side of (2.3). This completes the proof of our main result.

Next, we consider other variation of (2.1). In fact, we establish an integral formula for the generalized Bessel-Maitland function $J_{\eta, \lambda, \nu, \sigma, \delta, p}^{\mu, \rho, \gamma, q}(z)$, which is expressed in terms of the generalized hypergeometric function ${}_pF_q$.

Variation of (2.1): Let the conditions of our main result be satisfied, then the following integral formula holds true:

$$\begin{aligned} &\int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} J_{\eta, \lambda, \nu, \sigma, \delta, p}^{\mu, \rho, \gamma, q} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy \\ &= \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\lambda+1)\Gamma(\alpha+\beta)} \\ &\times {}_{\rho+q+3}F_{\sigma+p+\eta+2} \left[\begin{array}{c} \Delta(\rho, \mu), \Delta(q; \gamma), \alpha, \beta, \\ \Delta(\sigma, \nu), \Delta(p, \delta), \Delta(\eta; \lambda+1), \Delta(2; \alpha+\beta); \end{array} \begin{array}{c} 1, \\ ; \\ \frac{-a\rho^\rho q^q}{4\sigma^\sigma \eta^\eta p^p} \end{array} \right], \end{aligned} \quad (2.4)$$

where $\Delta(m; l)$ abbreviates the array of m parameters $\frac{l}{m}, \frac{l+1}{m}, \dots, \frac{l+m-1}{m}$, $m \geq 1$.

Proof: In order to prove the result (2.4), using the results

$$\Gamma(\alpha + n) = \Gamma(\alpha)(\alpha)_n$$

and

$$(l)_{kn} = k^{kn} \left(\frac{l}{k} \right)_n \left(\frac{l+1}{k} \right)_n \cdots \left(\frac{l+k-1}{k} \right)_n,$$

(Gauss multiplication theorem) in (2.3) and summing up the given series with the help of (1.16), we easily arrive at our required result (2.4).

3 Special Cases

In this section, we establish some special cases of generalized Bessel-Maitland functions as follows.

(i). On replacing λ by $\lambda - 1$ in (2.1) and then by using (1.11), we get:

$$\int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} E_{\eta, \lambda, \nu, \sigma, \delta, p}^{\mu, \rho, \gamma, q} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy$$

$$= \frac{\Gamma(\nu)\Gamma(\delta)}{\Gamma(\mu)\Gamma(\gamma)} {}_5\Psi_4 \left[\begin{array}{c} (\mu, \rho), (\gamma, q), (\alpha, 1), (\beta, 1), (1, 1); \\ (\nu, \sigma), (\delta, p), (\lambda, \eta), (\alpha + \beta, 2); \end{array} a \right], \quad (3.1)$$

where $\alpha, \beta, \eta, \gamma, \mu, \nu, \sigma, \rho, \delta, \lambda \in \mathbb{C}$, $p, q > 0$ and $q < \Re(\alpha) + p$.

(ii). On replacing λ by $\lambda - 1$ in (2.4) and then by using (1.11), we get:

$$\begin{aligned} & \int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} E_{\eta, \lambda, \nu, \sigma, \delta, p}^{\mu, \rho, \gamma, q} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy \\ &= \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\lambda)\Gamma(\alpha+\beta)} \\ & \times {}_{\rho+q+3}F_{\sigma+p+\eta+2} \left[\begin{array}{c} \Delta(\rho, \mu), \Delta(q; \gamma), \alpha, \beta, 1, ; \\ \Delta(\sigma, \nu), \Delta(p, \delta), \Delta(\eta; \lambda), \Delta(2; \alpha + \beta); \end{array} \frac{a\rho^\rho q^q}{4\sigma^\sigma \eta^\eta p^p} \right], \end{aligned} \quad (3.2)$$

where $\alpha, \beta, \eta, \gamma, \mu, \nu, \sigma, \rho, \delta, \lambda \in \mathbb{C}$, $p, q > 0$ and $q < \Re(\alpha) + p$.

(iii). On setting $\mu = \nu = \rho = \sigma = 1$ and replacing λ by $\lambda - 1$ in (2.1) and then by using (1.10), we find:

$$\begin{aligned} & \int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} E_{\eta, \lambda, p}^{\gamma, \delta, q} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy \\ &= \frac{\Gamma(\delta)}{\Gamma(\gamma)} {}_4\Psi_3 \left[\begin{array}{c} (\gamma, q), (\alpha, 1), (\beta, 1), (1, 1); \\ (\delta, p), (\lambda, \eta), (\alpha + \beta, 2); \end{array} a \right]. \end{aligned} \quad (3.3)$$

where $\alpha, \beta, \eta, \gamma, \delta, \lambda \in \mathbb{C}$, $p, q > 0$ and $q < \Re(\alpha) + p$.

(iv). On setting $\mu = \nu, \rho = \sigma$ and replacing λ by $\lambda - 1$ in (2.4) and then by using (1.10), we find:

$$\begin{aligned} & \int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} E_{\eta, \lambda, p}^{\gamma, \delta, q} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy \\ &= \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\lambda)\Gamma(\alpha+\beta)} \\ & \times {}_{q+3}F_{p+\eta+2} \left[\begin{array}{c} \Delta(q; \gamma), \alpha, \beta, 1, ; \\ \Delta(p, \delta), \Delta(\eta; \lambda), \Delta(2; \alpha + \beta); \end{array} a \right], \end{aligned} \quad (3.4)$$

where $\alpha, \beta, \eta, \gamma, \delta, \lambda \in \mathbb{C}$, $p, q > 0$ and $q < \Re(\alpha) + p$.

(v). On setting $\mu = \nu = \rho = \sigma = p = \delta = 1$ and replacing λ by $\lambda - 1$ in (2.1) and then by using (1.8), we find:

$$\begin{aligned} & \int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} E_{\eta, \lambda}^{\gamma, q} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy \\ &= \frac{1}{\Gamma(\gamma)} {}_2\Psi_1 \left[\begin{array}{c} (\gamma, q), (\alpha, 1), (\beta, 1), (1, 1); \\ (\lambda, \eta), (\alpha + \beta, 2); \end{array} a \right]. \end{aligned} \quad (3.5)$$

where $\alpha, \beta, \gamma, \eta, \lambda \in \mathbb{C}; \Re(\alpha) > 0, \Re(\beta) > 0, \Re(\gamma) > 0, \Re(\eta) > 0, \Re(\lambda) > 0, q \in (0, 1) \cup \mathbb{N}$.

(vi). On setting $\mu = \nu, \rho = \sigma, p = \delta = 1$ and replacing λ by $\lambda - 1$ in (2.4) and then by using (1.8), we find:

$$\begin{aligned} & \int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} E_{\eta,\lambda}^{\gamma,q} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy \\ &= \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\lambda)\Gamma(\alpha+\beta)} \\ & \times {}_{q+3}F_{\eta+2} \left[\begin{matrix} \Delta(q; \gamma), & \alpha, & \beta, & 1, & ; & \frac{aq^q}{4\eta^\eta} \\ \Delta(\eta; \lambda), & \Delta(2; \alpha+\beta); \end{matrix} \right], \end{aligned} \quad (3.6)$$

where $\alpha, \beta, \gamma, \eta, \lambda \in \mathbb{C}; \Re(\alpha) > 0, \Re(\beta) > 0, \Re(\gamma) > 0, \Re(\eta) > 0, \Re(\lambda) > 0, q \in (0, 1) \cup \mathbb{N}$.

(vii). On setting $\mu = \nu, \rho = \sigma, p = q = \delta = 1$ and replacing λ by $\lambda - 1$ in (2.1) and then by using (1.7), we obtain:

$$\begin{aligned} & \int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} E_{\eta,\lambda}^{\gamma} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy \\ &= \frac{1}{\Gamma(\gamma)} {}_4\Psi_2 \left[\begin{matrix} (\gamma, 1), (\alpha, 1), (\beta, 1), (1, 1); \\ (\lambda, \eta), (\alpha + \beta, 2); \end{matrix} a \right]. \end{aligned} \quad (3.7)$$

where $\alpha, \beta, \gamma, \eta, \lambda \in \mathbb{C}; \Re(\alpha) > 0, \Re(\beta) > 0, \Re(\gamma) > 0, \Re(\eta) > 0, \Re(\lambda) > 0$.

(viii). On setting $\mu = \nu, \rho = \sigma, p = q = \delta = 1$ and replacing λ by $\lambda - 1$ in (2.4) and then by using (1.8), we get:

$$\begin{aligned} & \int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} E_{\eta,\lambda}^{\gamma} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy \\ &= \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\lambda)\Gamma(\alpha+\beta)} \\ & \times {}_4F_{\eta+2} \left[\begin{matrix} \gamma, & \alpha, & \beta, & 1, & ; & \frac{a}{4\eta^\eta} \\ \Delta(\eta; \lambda), & \Delta(2; \alpha+\beta); \end{matrix} \right], \end{aligned} \quad (3.8)$$

where $\alpha, \beta, \gamma, \eta, \lambda \in \mathbb{C}; \Re(\alpha) > 0, \Re(\beta) > 0, \Re(\gamma) > 0, \Re(\eta) > 0, \Re(\lambda) > 0$.

(ix). On setting $\mu = \nu = \rho = \sigma = p = q = \delta = \gamma = 1$ and replacing λ by $\lambda - 1$ in (2.1) and then by using (1.6), we acquire:

$$\begin{aligned} & \int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} E_{\eta,\lambda} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy \\ &= {}_3\Psi_2 \left[\begin{matrix} (\alpha, 1), (\beta, 1), (1, 1); \\ (\lambda, \eta), (\alpha + \beta, 2); \end{matrix} a \right]. \end{aligned} \quad (3.9)$$

where $\alpha, \beta, \eta, \lambda \in \mathbb{C}; \Re(\alpha) > 0, \Re(\beta) > 0, \Re(\eta) > 0, \Re(\lambda) > 0..$

(x). On setting $\mu = \nu = \rho = \sigma = p = q = \delta = \gamma = 1$ and replacing λ by $\lambda - 1$ in (2.4) and then by using (1.6), we find:

$$\int_0^1 \int_0^1 y^\alpha (1-x)^{\alpha-1} (1-y)^{\beta-1} (1-xy)^{1-\alpha-\beta} E_{\eta,\lambda} \left[\frac{ay(1-x)(1-y)}{(1-xy)^2} \right] dx dy$$

$$\begin{aligned}
&= \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\lambda)\Gamma(\alpha+\beta)} \\
&\times {}_3F_{\eta+2} \left[\begin{array}{ccccc} \alpha, & \beta, & 1, & ; & \frac{a}{4\eta^\eta} \\ \Delta(\eta; \lambda), & \Delta(2; \alpha+\beta); & & & \end{array} \right], \tag{3.10}
\end{aligned}$$

where $\alpha, \beta, \eta, \lambda \in \mathbb{C}; \Re(\alpha) > 0, \Re(\beta) > 0, \Re(\eta) > 0, \Re(\lambda) > 0.$

References

- [1] Choi, J, Agarwal, P, Mathur, S and Purohit, S.D, Certain new integral formulas involving the generalized Bessel functions, Bull. Korean Math. Soc., 51(4)(2014), 995-1003.
- [2] Choi, J, Agarwal, P, Certain unified integrals involving a product of Bessel functions of first kind, Honam Mathematical J., 35(4)(2013), 667-677.
- [3] Choi, J, Agarwal, P, Certain unified integrals associated with Bessel functions, Boundary Value Problem, Vol 2013, 95, 2013.
- [4] Ghayasuddin, M, Khan, W. A, A new extension of Bessel Maitland function and its properties, Matematiki Vesnik, (2018), 1-11.
- [5] Edward, J, A treatise on the Integral calculus, Vol. II, Chelsea Publishing Company, New York, 1922.
- [6] Fox, C, The G and H-functions and symmetrical Fourier kernels, Trans. Amer. Math. Soc., 98(1961), 395-429.
- [7] Khan, W. A, Nisar, K S, Unified integral operator involving generalized Bessel-Maitland function, Proc. Jangjeon Mathe. Soc. 21(2018), 339 - 346.
- [8] Khan, W. A, Nisar, K S, Beta type integral formula associated with Wright generalized Bessel function, Acta Math. Univ. Comenianae, 1(2018), 117-125.
- [9] Khan, W. A, Nisar, K S, Choi, J, An integral formula of the Mellin transform type involving the extended Wright Bessel function, Far East J. Math., 102(2017), 2903-2912.
- [10] Khan, W. A, Nisar, K S, Ahmad, M, On certain integral transforms involving generalized Bessel-Maitland function, Submitted.
- [11] Khan, M. A, Ahmed, S, On some properties of the generalized Mittag-Leffler function, Springer Plus 2013, 2:337.
- [12] Mittag-Leffler, G. M, Sur la nouvelle fonction $E_\alpha(x)$, C. R. Acad. Sci. Paris, 137(1903), 554-558.
- [13] Marichev, O. I, Handbook of integral transform and Higher transcendental functions. Theory and algorithm tables, Ellis Horwood, Chichester, New York, 1983.
- [14] Prabhakar, T. R, A singular integral equation with a generalized Mittag-Leffler function in the kernel, Yokohama Math. J., 19(1971), 715.
- [15] Rainville, E. D, Special functions, Macmillan, New York, 1960.
- [16] Samko, S. G, Kilbas, A.A and Marichev, O.I, Fractional Integrals and derivatives. Theory and Applications, Gordon and Breach, Yverdon et al., 1993.
- [17] Salim, T. O and Faraj, W, A generalization of Mittag-Leffler function and integral operator associated with fractional calculus, J. Frac. Calc. Appl., 3(5)(2012), 1-13.
- [18] Salim, T. O, Some properties relating to generalized Mittag-Leffler function, Adv. Appl. Math. Anal., 4(1)(2009), 21-30.
- [19] Shukla, A. K and Prajapati, J.C, On a generalization of Mittag-Leffler function and its properties, J. Math. Anal. Appl., 336(2007), 797-811.
- [20] Singh, M, Khan, M. A, Khan, A. H, On some properties of a generalization of Bessel-Maitland function, Int. J. Math. Trends and Tech., 14(1) (2014), 46-54.
- [21] Watson, G. N, A treatise on the theory of Bessel functions, Cambridge University Press, 1962.
- [22] Wiman, A, Uber den fundamental Satz in der Theorie der Funktionen $E_\alpha(x)$, Acta Math., 29(1905), 191-201.
- [23] Wright, E. M, On the coefficients of power series having exponential singularities., Proc. London Math. Soc., 8(1933), 71-79.
- [24] Wright, E. M, The asymptotic expansion of the generalized hypergeometric function, J. London Math. Soc., 10(1935), 286-293.

Author information

Musharraf Ali, (Corresponding Author), Department of Mathematics, G.F. College, Shahjahanpur-242001, India.

E-mail: drmusharrafali@gmail.com

Waseem A. Khan, Department of Mathematics and Natural Sciences Prince Mohammad Bin Fahd University, P.O Box 1664, Al Khobar 31952, Saudi Arabia.

E-mail: waseem08_khan@rediffmail.com

Idrees A. Khan, Department of Mathematics, Faculty of Science, Integral University, Lucknow-226026, India.

E-mail: khanidrees077@gmail.com

Received: August 2, 2018.

Accepted: November 22, 2018.