

# A study on two coupled modified KdV systems with time-dependent and constant coefficients

Abdul-Majid Wazwaz

Communicated by Ayman Badawi

2010 Mathematics Subject Classification: 37K25, 37K10, 35Q53

Keywords: Coupled mKdV equation; bright solitons; multiple soliton solutions; bilinear method; resonance.

## 1 Abstract

This work concerns itself on two new coupled mKdV systems, the first with time-dependent coefficients, and the second with constant coefficients. The soliton ansatz will be used for the first system to obtain 1-bright soliton solution. The simplified form of the bilinear method will be used to derive multiple-soliton solutions and multiple singular soliton solutions for the second coupled mKdV system.

## 2 Introduction

In solitary waves theory, the integrable coupled mKdV systems attract the interest of many mathematicians and physicists. The coupled variable coefficient mKdV is usually derived from a two-layer fluid model by using the multiple scale scheme with the reductive perturbation method.

Various methods [1–19] have been used to conduct analysis on the coupled nonlinear KdV and mKdV equations. Examples of the methods that have been used so far are the Hirota bilinear method, the Bäcklund transformation method, Darboux transformation, Pfaffian technique, the inverse scattering method, the Painlevé analysis, the generalized symmetry method, the subsidiary ordinary differential equation method (sub-ODE for short), the coupled amplitude-phase formulation, sine-cosine method, sech-tanh method, the mapping and the deformation approach, and many other methods. The Hirota's bilinear method [7–14], and the Hereman's simplified form [15] are rather heuristic and significant. These approaches possess powerful features that make it practical for the determination of multiple soliton solutions [16–31] for a wide class of nonlinear evolution equations. Moreover, the soliton ansatz [3] has been used for the determination of the bright soliton solutions.

In this work, two coupled modified KdV (mKdV) equations, the first with time-dependent coefficients, and the second with constant coefficients, will be investigated for the determination of multiple soliton solutions and bright soliton solutions. The coupled mKdV equations that we will examine are given by

$$\begin{aligned}u_t + f(t)u_{xxx} + g(t)u^2u_x + h(t)uvw_x &= 0, \\v_t + f(t)v_{xxx} + g(t)v^2u_x + h(t)vwu_x &= 0, \\w_t + f(t)w_{xxx} + g(t)w^2w_x + h(t)wuv_x &= 0,\end{aligned}\tag{1}$$

where  $f(t)$ ,  $g(t)$  and  $h(t)$  are time-dependent coefficients. If we set  $f(t) = 1$ ,  $g(t) = 6$ , and  $h(t) = 3$ , Eq. (1) will be reduced to

$$\begin{aligned}u_t + u_{xxx} + 6u^2u_x + 3uvw_x &= 0, \\v_t + v_{xxx} + 6v^2u_x + 3vwu_x &= 0, \\w_t + w_{xxx} + 6w^2w_x + 3wuv_x &= 0.\end{aligned}\tag{2}$$

The first coupled KdV equation (1) will be studied for bright solitons, whereas the latter will be investigated for complete integrability and for the determination of multiple soliton solutions and multiple singular soliton solutions. The computer algebra systems such as Maple and Mathematica allow us to perform complicated and tedious calculations.

This work is organized as follows. In Section 2, we make use of the soliton ansatz to determine bright solitons for the first model of coupled mKdV equations with time-dependent coupling coefficients. In Section 3, we derive multiple soliton solutions and multiple singular soliton solutions for the second model of coupled mKdV equations with constant linear coupling coefficients. Conclusions are contained in Section 4.

### 3 The first coupled mKdV system

We first begin our analysis on the first coupled mKdV system with time-dependent coefficients

$$\begin{aligned} u_t + f(t)u_{xxx} + g(t)u^2u_x + h(t)uvw_x &= 0, \\ v_t + f(t)v_{xxx} + g(t)v^2u_x + h(t)vwu_x &= 0, \\ w_t + f(t)w_{xxx} + g(t)w^2w_x + h(t)wuv_x &= 0, \end{aligned} \quad (1)$$

where  $u(x, t)$ ,  $v(x, t)$  and  $w(x, t)$  are real functions of the spatial variable  $x$  and the temporal variable  $t$ ,  $f(t)$ ,  $g(t)$  and  $h(t)$  are real coupling coefficients which are time-dependent functions, and subscripts denote partial derivatives.

As stated before, this model is derived from a two-layer fluid model which is used to study the interaction between the atmosphere and oceanic phenomena. The derivation of (1) was achieved by using the multiple scale approach with the reductive perturbation method [1].

In this section, our goal is to find the exact bright soliton solutions for the coupled mKdV system (1), governing the mutual interaction in addition to nonlinear self-interaction of two solitary waves in a mKdV system. For definiteness, we look to study the problem when linear and nonlinear coupling terms are present. It should be mentioned that consideration of linear coupling terms to study the propagation of soliton pulses is just a particular case because most of real physical systems exhibit nonlinear interaction between waves.

To obtain the bright soliton solutions of (1), we assume the solitary wave ansätze of the form [3–5]

$$\begin{aligned} u(x, t) &= A(t)\operatorname{sech}^p \{ \eta(t)(x - vt) \}, \\ v(x, t) &= B(t)\operatorname{sech}^q \{ \eta(t)(x - vt) \}, \\ w(x, t) &= C(t)\operatorname{sech}^r \{ \eta(t)(x - vt) \}, \end{aligned} \quad (2)$$

where  $v = v(t)$  is a time-dependent coefficient that will be determined. Here  $A$ ,  $B$ , or  $C$  is the amplitude of solitons,  $\eta$  and  $v$  are the inverse width and the velocity of the solitons respectively. The exponents  $p$ ,  $q$  and  $r$  are unknown at this point and will be determined later.

From the first part of ansatz (2), one obtains

$$u_t = \frac{dA}{dt}\operatorname{sech}^p\theta - Ap \left\{ \frac{d\eta}{dt}(x - vt) - \eta \left( v + t \frac{dv}{dt} \right) \right\} \operatorname{sech}^p\theta \tanh\theta, \quad (3)$$

$$u_x = -A p \eta \operatorname{sech}^p\theta \tanh\theta, \quad (4)$$

$$u_{xxx} = -A (p\eta)^3 \operatorname{sech}^p\theta \tanh\theta + p(p+1)(p+2)\eta^3 A \operatorname{sech}^{p+2}\theta \tanh\theta. \quad (5)$$

From the second part of ansatz (2) we find

$$v_t = \frac{dB}{dt} \operatorname{sech}^q \theta - Bq \left\{ \frac{d\eta}{dt} (x - vt) - \eta \left( v + t \frac{dv}{dt} \right) \right\} \operatorname{sech}^q \theta \tanh \theta, \quad (6)$$

$$v_x = -Bq\eta \operatorname{sech}^q \theta \tanh \theta, \quad (7)$$

$$v_{xxx} = -B(q\eta)^3 \operatorname{sech}^q \theta \tanh \theta + q(q+1)(q+2)\eta^3 B \operatorname{sech}^{q+2} \theta \tanh \theta. \quad (8)$$

From the third part of ansatz (2) we find

$$w_t = \frac{dC}{dt} \operatorname{sech}^r \theta - Cr \left\{ \frac{d\eta}{dt} (x - vt) - \eta \left( v + t \frac{dv}{dt} \right) \right\} \operatorname{sech}^r \theta \tanh \theta, \quad (9)$$

$$w_x = -Cr\eta \operatorname{sech}^r \theta \tanh \theta, \quad (10)$$

$$w_{xxx} = -C(r\eta)^3 \operatorname{sech}^r \theta \tanh \theta + r(r+1)(r+2)\eta^3 C \operatorname{sech}^{r+2} \theta \tanh \theta, \quad (11)$$

where

$$\theta = \eta(t) (x - v(t)t). \quad (12)$$

Substituting Eqs. (2)–(12) into Eqs. (1) gives

$$\begin{aligned} & \operatorname{sech}^p \theta \frac{dA}{dt} - pA \left\{ \frac{d\eta}{dt} (x - vt) - \eta \left( v + t \frac{dv}{dt} \right) \right\} \operatorname{sech}^p \theta \tanh \theta \\ & + f(t) \left\{ -A(p\eta)^3 \operatorname{sech}^p \theta \tanh \theta + p(p+1)(p+2)\eta^3 A \operatorname{sech}^{p+2} \theta \tanh \theta \right\} \\ & + g(t) \left\{ -A^3 p\eta \operatorname{sech}^{3p} \theta \tanh \theta \right\} \\ & + h(t) \left\{ -ABC r\eta \operatorname{sech}^{p+q+r} \theta \tanh \theta \right\} = 0, \end{aligned} \quad (13)$$

$$\begin{aligned} & \operatorname{sech}^q \theta \frac{dB}{dt} - qB \left\{ \frac{d\eta}{dt} (x - vt) - \eta \left( v + t \frac{dv}{dt} \right) \right\} \operatorname{sech}^q \theta \tanh \theta \\ & + f(t) \left\{ -B(q\eta)^3 \operatorname{sech}^q \theta \tanh \theta + q(q+1)(q+2)\eta^3 B \operatorname{sech}^{q+2} \theta \tanh \theta \right\} \\ & + g(t) \left\{ -AB^2 p\eta \operatorname{sech}^{2p+q} \theta \tanh \theta \right\} \\ & + h(t) \left\{ -ABC p\eta \operatorname{sech}^{p+q+r} \theta \tanh \theta \right\} = 0, \end{aligned} \quad (14)$$

and

$$\begin{aligned} & \operatorname{sech}^r \theta \frac{dC}{dt} - rC \left\{ \frac{d\eta}{dt} (x - vt) - \eta \left( v + t \frac{dv}{dt} \right) \right\} \operatorname{sech}^r \theta \tanh \theta \\ & + f(t) \left\{ -C(r\eta)^3 \operatorname{sech}^r \theta \tanh \theta + r(r+1)(r+2)\eta^3 C \operatorname{sech}^{r+2} \theta \tanh \theta \right\} \\ & + g(t) \left\{ -C^3 r\eta \operatorname{sech}^{3r} \theta \tanh \theta \right\} \\ & + h(t) \left\{ -ABC q\eta \operatorname{sech}^{p+q+r} \theta \tanh \theta \right\} = 0. \end{aligned} \quad (15)$$

Equating the exponents of  $\operatorname{sech}^{p+2} \theta \tanh \theta$  and  $\operatorname{sech}^{3p} \theta \tanh \theta$  terms in Eq. (13), one gets

$$p + 2 = 3p, \quad (16)$$

so that

$$p = 1. \quad (17)$$

We next match the exponents of  $\text{sech}^{q+2}\theta \tanh \theta$  and  $\text{sech}^{p+2q}\theta \tanh \theta$  in (14) to obtain

$$q + 2 = p + 2q, \quad (18)$$

which gives

$$q = 1. \quad (19)$$

We finally match  $\text{sech}^{r+2}\theta \tanh \theta$  and  $\text{sech}^{3r}\theta \tanh \theta$  in (15) to obtain

$$r + 2 = 3r, \quad (20)$$

which gives

$$r = 1. \quad (21)$$

If we put  $p = 1$  in Eq. (13), we can determine the inverse width  $\eta$  of the soliton pulses by setting the corresponding coefficients of  $\text{sech}^3\theta \tanh \theta$  to zero, hence we find

$$6Af(t)\eta^3 - A^3g(t)\eta - ABCh(t)\eta = 0, \quad (22)$$

which gives

$$\eta = \sqrt{\frac{A^2g(t) + BCh(t)}{6f(t)}}, f(t) \neq 0. \quad (23)$$

Similarly we find from setting the coefficients of  $\text{sech}^3\theta \tanh \theta$  terms to zero in Eq. (14) that

$$\eta = \sqrt{\frac{ABg(t) + ACh(t)}{6f(t)}}, f(t) \neq 0. \quad (24)$$

Proceeding as before, we find from (15) that

$$\eta = \sqrt{\frac{C^2g(t) + ABh(t)}{6f(t)}}, f(t) \neq 0. \quad (25)$$

Equating the three values of  $\eta$  from (23)–(25) gives

$$A(t) = B(t) = C(t). \quad (26)$$

This in turn gives the inverse width by

$$\eta = A(t) \sqrt{\frac{g(t) + h(t)}{6f(t)}}, f(t) \neq 0. \quad (27)$$

Setting the coefficients of  $\text{sech}^p\theta \tanh \theta$ ,  $\text{sech}^q\theta \tanh \theta$ , and  $\text{sech}^r\theta \tanh \theta$  terms to zero in Eqs. (13) – (15), respectively, we get

$$-pA \left\{ \frac{d\eta}{dt} (x - vt) - \eta \left( v + t \frac{dv}{dt} \right) \right\} - f(t)A(p\eta)^3 = 0, \quad (28)$$

$$-qB \left\{ \frac{d\eta}{dt} (x - vt) - \eta \left( v + t \frac{dv}{dt} \right) \right\} - f(t)B(q\eta)^3 = 0, \quad (29)$$

and

$$-rC \left\{ \frac{d\eta}{dt} (x - vt) - \eta \left( v + t \frac{dv}{dt} \right) \right\} - f(t)C(r\eta)^3 = 0. \quad (30)$$

Taking into account the fact that the soliton velocity  $v(t)$  we want to determine from Eqs. (28) – (30) is a function of time, one can split these three equations into four equations as follows:

$$\frac{d\eta}{dt} = 0, \quad (31)$$

$$\frac{d(vt\eta)}{dt} - f\eta^3 = 0, \quad (32)$$

$$\frac{d(vt\eta)}{dt} - f\eta^3 = 0, \quad (33)$$

$$\frac{d(vt\eta)}{dt} - f\eta^3 = 0. \quad (34)$$

It is clear that (32)–(34) are equivalent. By integrating the above equations with respect to the time variable  $t$ , one obtains

$$\eta(t) = k, \quad (35)$$

and

$$v(t) = \frac{k^2}{t} \int_0^t f(t') dt', \quad (36)$$

where  $k$  is an integral constant related to the initial pulse inverse width as defined in (35). From Eq. (35), it is apparent that the inverse width of the solitons remains constant when the pulse propagates in the varying mKdV system. We remark also from Eq. (36) that the pulse velocity is affected by the time-dependent coupling coefficients  $f(t)$ .

Lastly, we find from setting the coefficients of  $\text{sech}^p\theta$  and  $\text{sech}^q\theta$  terms to zero in Eqs. (13)–(15), respectively, that

$$\frac{dA}{dt} = 0, \quad (37)$$

$$\frac{dB}{dt} = 0, \quad (38)$$

$$\frac{dC}{dt} = 0, \quad (39)$$

which gives after integration

$$A(t) = A_0, \quad (40)$$

$$B(t) = B_0, \quad (41)$$

$$C(t) = C_0, \quad (42)$$

where  $A_0, B_0$  and  $C_0$  are integral constants related to the initial pulse amplitudes of the three solitons. As  $A(t) = B(t) = C(t)$  from (26), we get  $A_0 = B_0 = C_0 = \alpha$  where  $\alpha$  is a constant.

It is interesting to note also from (26) that, since  $A(t) = \alpha$ , and the inverse width  $\eta = k$ , then it is necessary that

$$\frac{g(t) + h(t)}{6f(t)} = \beta. \quad (43)$$

Finally we find that the three bright soliton solutions are symmetric and given by

$$u(x, t) = v(x, t) = w(x, t) = A_0 \text{sech} \{k(x - v(t)t)\}, \quad (44)$$

where the velocity  $v(t)$  is given by (36) and  $k$  is given by (35).

One concludes that the first coupled equation (1) possesses the soliton solution and anti-soliton solutions since  $u(x, t) = \pm v(x, t)$ .

#### 4 The second coupled mKdV system

In this section, we will study the second coupled mKdV system, with constant coefficients, given by

$$\begin{aligned} u_t + u_{xxx} + 6u^2u_x + 3uvw_x &= 0, \\ v_t + v_{xxx} + 6v^2v_x + 3vuw_x &= 0, \\ w_t + w_{xxx} + 6w^2w_x + 3wuv_x &= 0. \end{aligned} \quad (1)$$

##### 4.1 Multiple soliton solutions

Substituting

$$\begin{aligned} u(x, t) &= e^{\theta_i}, \theta_i = k_i x - c_i t, \\ v(x, t) &= A e^{\theta_i}, \\ w(x, t) &= B e^{\theta_i}, \end{aligned} \quad (2)$$

where  $A$  and  $B$  are constants, into the linear terms of (1) gives the dispersion relation by

$$c_i = k_i^3, \quad (3)$$

and as a result we obtain

$$\theta_i = k_i x - k_i^3 t. \quad (4)$$

The multi-soliton solutions of the coupled mKdV system are given by

$$\begin{aligned} u(x, t) &= R \left( \arctan\left(\frac{f(x, t)}{g(x, t)}\right) \right)_x = R \frac{f_x g - g_x f}{f^2 + g^2}, \\ v(x, t) &= R_1 \left( \arctan\left(\frac{f(x, t)}{g(x, t)}\right) \right)_x = R_1 \frac{f_x g - g_x f}{f^2 + g^2}, \\ w(x, t) &= R_2 \left( \arctan\left(\frac{f(x, t)}{g(x, t)}\right) \right)_x = R_2 \frac{f_x g - g_x f}{f^2 + g^2}, \end{aligned} \quad (5)$$

where the auxiliary functions  $f(x, t)$  and  $g(x, t)$  for the single soliton solution are given by

$$\begin{aligned} f(x, t) &= 1 + e^{\theta_1} = 1 + e^{k_1 x - k_1^3 t}, \\ g(x, t) &= 1 - e^{\theta_1} = 1 - e^{k_1 x - k_1^3 t}. \end{aligned} \quad (6)$$

Substituting (5) and (6) into (1) and solving for  $R, R_1$  and  $R_2$  we find three sets of solutions given by

$$R = \frac{2}{\sqrt{7}}, R_1 = \frac{12}{\sqrt{7}}, R_2 = \frac{4}{\sqrt{7}}, \quad (7)$$

$$R = -\frac{4}{\sqrt{3}}, R_1 = -\frac{4}{\sqrt{3}}, R_2 = \frac{2}{\sqrt{3}}, \quad (8)$$

and

$$R = 2\sqrt{\frac{2}{3}}, R_1 = 2\sqrt{\frac{2}{3}}, R_2 = 2\sqrt{\frac{2}{3}}. \quad (9)$$

Combining (5)–(9) gives the following three sets of single soliton solutions

$$\begin{aligned} u(x, t) &= \frac{2k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{7}(1+e^{2k_1(x-(k_1^2+\alpha)t})}), \\ v(x, t) &= \frac{12k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{7}(1+e^{2k_1(x-(k_1^2+\alpha)t})}), \\ w(x, t) &= \frac{4k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{7}(1+e^{2k_1(x-(k_1^2+\alpha)t})}), \end{aligned} \quad (10)$$

$$\begin{aligned} u(x, t) &= -\frac{4k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{3}(1+e^{2k_1(x-(k_1^2+\alpha)t})}), \\ v(x, t) &= -\frac{4k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{3}(1+e^{2k_1(x-(k_1^2+\alpha)t})}), \\ w(x, t) &= \frac{2k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{3}(1+e^{2k_1(x-(k_1^2+\alpha)t})}), \end{aligned} \quad (11)$$

and

$$\begin{aligned} u(x, t) &= \frac{2\sqrt{2}k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{3}(1+e^{2k_1(x-(k_1^2+\alpha)t})}), \\ v(x, t) &= \frac{2\sqrt{2}k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{3}(1+e^{2k_1(x-(k_1^2+\alpha)t})}), \\ w(x, t) &= \frac{2\sqrt{2}k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{3}(1+e^{2k_1(x-(k_1^2+\alpha)t})}). \end{aligned} \quad (12)$$

The last set gives symmetric solutions similar to what we obtained before for coupled mKdV system with time-dependent coefficients.

To determine the two-soliton solutions we set

$$\begin{aligned} f(x, t) &= 1 + e^{\theta_1} + e^{\theta_2} - a_{12}e^{\theta_1+\theta_2}, \\ g(x, t) &= 1 - e^{\theta_1} - e^{\theta_2} - a_{12}e^{\theta_1+\theta_2}. \end{aligned} \quad (13)$$

Substituting (13) into (5) and using the obtained result in the coupled mKdV system (1), one obtains the phase shift  $a_{12}$  by

$$a_{12} = \frac{(k_1 - k_2)^2}{(k_1 + k_2)^2}, \quad (14)$$

and this can be generalized to

$$a_{ij} = \frac{(k_i - k_j)^2}{(k_i + k_j)^2}, \quad 1 \leq i < j \leq 3. \quad (15)$$

Substituting (14) and (13) into (5), and using the three sets (7)–(9) we obtain three sets of two solitons solutions. It is interesting to point out that the system (1) does not show any resonant phenomenon [8] because the phase shift term  $a_{12}$  in (14) cannot be 0 or  $\infty$  for  $|k_1| \neq |k_2|$ .

To determine the three soliton solutions, we set

$$\begin{aligned} f(x, t) &= 1 + e^{\theta_1} + e^{\theta_2} + e^{\theta_3} \\ &\quad - a_{12}e^{\theta_1+\theta_2} - a_{13}e^{\theta_1+\theta_3} - a_{23}e^{\theta_2+\theta_3} - b_{123}e^{\theta_1+\theta_2+\theta_3}, \\ g(x, t) &= 1 - e^{\theta_1} - e^{\theta_2} - e^{\theta_3} \\ &\quad - a_{12}e^{\theta_1+\theta_2} - a_{13}e^{\theta_1+\theta_3} - a_{23}e^{\theta_2+\theta_3} + b_{123}e^{\theta_1+\theta_2+\theta_3}, \end{aligned} \quad (16)$$

where the phase shifts  $a_{ij}$  are derived above in (15). Proceeding as before we find

$$b_{123} = a_{12}a_{13}a_{23}. \quad (17)$$

This shows that the coupled mKdV system (1) is completely integrable and  $N$ -soliton solutions can be obtained for finite  $N$ , where  $N \geq 1$ . Three sets of three soliton solutions for the coupled mKdV equation (1) are therefore obtained by substituting (16) into (5).

## 4.2 Multiple singular soliton solutions

In this section, we will determine multiple singular soliton solutions for the coupled mKdV system (1). Following [13–16], the singular soliton solution of the coupled mKdV system (1) is assumed to be of the form

$$\begin{aligned} u(x, t) &= R \left( \ln \left( \frac{f(x, t)}{g(x, t)} \right) \right)_x = R \frac{gf_x - fg_x}{gf}, \\ v(x, t) &= R_1 \left( \ln \left( \frac{f(x, t)}{g(x, t)} \right) \right)_x = R_1 \frac{gf_x - fg_x}{gf}, \\ w(x, t) &= R_2 \left( \ln \left( \frac{f(x, t)}{g(x, t)} \right) \right)_x = R_2 \frac{gf_x - fg_x}{gf}, \end{aligned} \quad (18)$$

where  $R, R_1$  and  $R_2$  are constants that will be determined. The auxiliary functions  $f(x, t)$  and  $g(x, t)$  have expansions of the form

$$\begin{aligned} f(x, t) &= 1 + \sum_{n=1}^{\infty} f_n(x, t), \\ g(x, t) &= 1 - \sum_{n=1}^{\infty} g_n(x, t). \end{aligned} \quad (19)$$

Following the discussion presented in the previous section, the dispersion relation is given by

$$c_i = k_i^3, \quad (20)$$

and as a result we obtain

$$\theta_i = k_i x - k_i^3 t. \quad (21)$$

The obtained results give a new definition to (19) in the form

$$\begin{aligned} f(x, t) &= 1 + e^{k_1(x - k_1^2 t)}, \\ g(x, t) &= 1 - e^{k_1(x - k_1^2 t)}. \end{aligned} \quad (22)$$

Substituting (22) into (18), and using the outcome in (1), one obtains three sets of solutions given by

$$R = \frac{1}{\sqrt{7}}, R_1 = \frac{6}{\sqrt{7}}, R_2 = \frac{2}{\sqrt{7}}, \quad (23)$$

$$R = -\frac{2}{\sqrt{3}}, R_1 = -\frac{2}{\sqrt{3}}, R_2 = \frac{1}{\sqrt{3}}, \quad (24)$$

and

$$R = \sqrt{\frac{2}{3}}, R_1 = \sqrt{\frac{2}{3}}, R_2 = \sqrt{\frac{2}{3}}. \quad (25)$$

Combining the previous results gives the following three sets of single singular soliton solutions

$$\begin{aligned} u(x, t) &= \frac{2k_1 e^{k_1(x - (k_1^2 + \alpha)t)}}{\sqrt{7}(1 - e^{2k_1(x - (k_1^2 + \alpha)t)}}), \\ v(x, t) &= \frac{12k_1 e^{k_1(x - (k_1^2 + \alpha)t)}}{\sqrt{7}(1 - e^{2k_1(x - (k_1^2 + \alpha)t)}}), \\ w(x, t) &= \frac{4k_1 e^{k_1(x - (k_1^2 + \alpha)t)}}{\sqrt{7}(1 - e^{2k_1(x - (k_1^2 + \alpha)t)}}), \end{aligned} \quad (26)$$

$$\begin{aligned} u(x, t) &= -\frac{4k_1 e^{k_1(x - (k_1^2 + \alpha)t)}}{\sqrt{3}(1 - e^{2k_1(x - (k_1^2 + \alpha)t)}}), \\ v(x, t) &= -\frac{4k_1 e^{k_1(x - (k_1^2 + \alpha)t)}}{\sqrt{3}(1 - e^{2k_1(x - (k_1^2 + \alpha)t)}}), \\ w(x, t) &= \frac{2k_1 e^{k_1(x - (k_1^2 + \alpha)t)}}{\sqrt{3}(1 - e^{2k_1(x - (k_1^2 + \alpha)t)}}), \end{aligned} \quad (27)$$

and

$$\begin{aligned} u(x, t) &= \frac{2\sqrt{2}k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{3}(1-e^{2k_1(x-(k_1^2+\alpha)t})}), \\ v(x, t) &= \frac{2\sqrt{2}k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{3}(1-e^{2k_1(x-(k_1^2+\alpha)t})}), \\ w(x, t) &= \frac{2\sqrt{2}k_1 e^{k_1(x-(k_1^2+\alpha)t)}}{\sqrt{3}(1-e^{2k_1(x-(k_1^2+\alpha)t})}). \end{aligned} \quad (28)$$

The last set gives symmetric solutions. The singularity behavior of each solution is clearly observed from the denominators.

To determine the singular two soliton solutions, we set

$$\begin{aligned} f(x, t) &= 1 + e^{\theta_1} + e^{\theta_2} + a_{12}e^{\theta_1+\theta_2}, \\ g(x, t) &= 1 - e^{\theta_1} - e^{\theta_2} + b_{12}e^{\theta_1+\theta_2}. \end{aligned} \quad (29)$$

Substituting (29) into (19) and using the outcome into (1), we find that (29) is a solution of this equation if the phase shifts  $a_{12}$  and  $b_{12}$ , and therefore  $a_{ij}$  and  $b_{ij}$ , are equal and given by

$$a_{ij} = b_{ij} = \frac{(k_i - k_j)^2}{(k_i + k_j)^2}. \quad (30)$$

For the singular two soliton solutions we use  $1 \leq i < j \leq 2$  to obtain

$$\begin{aligned} f(x, t) &= 1 + e^{k_1(x-k_1^2t)} + e^{k_2(x-k_2^2t)} + \frac{(k_1-k_2)^2}{(k_1+k_2)^2} e^{(k_1+k_2)x-(k_1^3+k_2^3)t}, \\ g(x, t) &= 1 - e^{k_1(x-k_1^2t)} - e^{k_2(x-k_2^2t)} + \frac{(k_1-k_2)^2}{(k_1+k_2)^2} e^{(k_1+k_2)x-(k_1^3+k_2^3)t}. \end{aligned} \quad (31)$$

This in turn gives the singular two soliton solutions if we substitute (31) into (18).

To determine the singular three soliton solutions, we can proceed in a similar manner and set

$$\begin{aligned} f(x, t) &= 1 + e^{\theta_1} + e^{\theta_2} + e^{\theta_3} \\ &+ a_{12}e^{\theta_1+\theta_2} + a_{23}e^{\theta_2+\theta_3} + a_{13}e^{\theta_1+\theta_3} + f_3(x, t), \\ g(x, t) &= 1 - e^{\theta_1} - e^{\theta_2} - e^{\theta_3} \\ &+ a_{12}e^{\theta_1+\theta_2} + a_{23}e^{\theta_2+\theta_3} + a_{13}e^{\theta_1+\theta_3} + g_3(x, t). \end{aligned} \quad (32)$$

Substituting (32) into (18) and using the result into (1) to find that

$$\begin{aligned} f_3(x, t) &= b_{123}e^{\theta_1+\theta_2+\theta_3}, \\ g_3(x, t) &= -b_{123}e^{\theta_1+\theta_2+\theta_3}, \\ b_{123} &= a_{12}a_{13}a_{23}. \end{aligned} \quad (33)$$

For the singular three-soliton solutions we use  $1 \leq i < j \leq 3$ , we therefore obtain

$$\begin{aligned} f(x, t) &= 1 + e^{k_1(x-k_1^2t)} + e^{k_2(x-k_2^2t)} + e^{k_3(x-k_3^2t)} \\ &+ \frac{(k_1-k_2)^2}{(k_1+k_2)^2} e^{(k_1+k_2)x-(k_1^3+k_2^3)t} + \frac{(k_1-k_3)^2}{(k_1+k_3)^2} e^{(k_1+k_3)x-(k_1^3+k_3^3)t} \\ &+ \frac{(k_2-k_3)^2}{(k_2+k_3)^2} e^{(k_2+k_3)x-(k_2^3+k_3^3)t} \\ &+ \frac{(k_1-k_2)^2(k_1-k_3)^2(k_2-k_3)^2}{(k_1+k_2)^2(k_1+k_3)^2(k_2+k_3)^2} e^{(k_1+k_2+k_3)x-(k_1^3+k_2^3+k_3^3)t}, \\ g(x, t) &= 1 - e^{k_1(x-k_1^2t)} - e^{k_2(x-k_2^2t)} - e^{k_3(x-k_3^2t)} \\ &+ \frac{(k_1-k_2)^2}{(k_1+k_2)^2} e^{(k_1+k_2)x-(k_1^3+k_2^3)t} + \frac{(k_1-k_3)^2}{(k_1+k_3)^2} e^{(k_1+k_3)x-(k_1^3+k_3^3)t} \\ &+ \frac{(k_2-k_3)^2}{(k_2+k_3)^2} e^{(k_2+k_3)x-(k_2^3+k_3^3)t} \\ &- \frac{(k_1-k_2)^2(k_1-k_3)^2(k_2-k_3)^2}{(k_1+k_2)^2(k_1+k_3)^2(k_2+k_3)^2} e^{(k_1+k_2+k_3)x-(k_1^3+k_2^3+k_3^3)t}. \end{aligned} \quad (34)$$

The singular three-soliton solutions follow immediately upon substituting (34) into (18).

## 5 Discussion

In this work we studied two coupled mKdV systems. The first system includes time-dependent coefficients, where we obtained only symmetric solutions with the wave speed  $v(t)$  depends on the function  $f(t)$ . The second model with constant coefficients was handled by the simplified bilinear method. Three sets of multiple soliton solutions were obtained, one set consists of symmetric solutions whereas the other two sets are asymmetric solutions. Moreover, three sets of multiple singular soliton solutions, one of them is symmetric, were obtained. The approach used for the second system is different than the approach used by Hietarinta [14]. We believe that this approach will be applicable to other applications.

## References

- [1] Y. J.-Rong and M. J.-Jian, Complexiton solutions of a special coupled mKdV equation, *Chin. Phys. Lett.*, 25 (2008) 1527–1530.
- [2] H. Triki, A.M. Wazwaz, Sub-ODE method and soliton solutions for the variable- coefficient mKdV equation, *Appl. Math. Comput.* 214 (2009) 370–373.
- [3] H. Triki, A.M. Wazwaz, Bright and dark soliton solutions for a  $K(m, n)$  equation with  $t$ -dependent coefficients, *Phys. Lett A* 373 (2009) 2162–215.
- [4] A. Biswas, 1-soliton solution of the B(m, n) equation with generalized evolution, *Commun. Nonlinear Sci. Numer. Simul.* 14 (2009) 3226–3229.
- [5] A. Biswas, 1-soliton solution of the K(m, n) equation with generalized evolution, *Phys. Lett. A* 372 (2008) 4601–4602.
- [6] M. Dehghan and A. Shokri, A numerical method for solution of the two-dimensional sine-Gordon equation using the radial basis functions, *Comput. Math. Simulation*, 79 (2008) 700–715.
- [7] R.Hirota, Discretization of coupled modified KdV equations, *Chaos, Solitons and Fractals*, 11 (2000) 77–84.
- [8] R. Hirota and M. Ito, Resonance of solitons in one dimension, *J. Phys. Soc. Japan*, 52(3), (1983) 744–748.
- [9] R. Hirota and J.Satsuma,  $N$ -soliton solutions of model equations for shallow water waves, *J. Phys. Soc. Japan*, 40(2) (1976) 611–612.
- [10] R. Hirota, A new form of Bäcklund transformations and its relation to the inverse scattering problem, *Progress of Theoretical Physics*, 52(5), (1974) 1498–1512.
- [11] R. Hirota, *The Direct Method in Soliton Theory*, Cambridge University Press, Cambridge (2004).
- [12] R. Hirota, Exact solutions of the Korteweg-de Vries equation for multiple collisions of solitons, *Physical Review Letters*, 27(18) (1971) 1192–1194.
- [13] J. Hietarinta, A search for bilinear equations passing Hirota's three-soliton condition. I. KdV-type bilinear equations, *J. Math. Phys.*, 28(8)(1987) 1732–1742.
- [14] J. Hietarinta, A search for bilinear equations passing Hirota's three-soliton condition. II. mKdV-type bilinear equations, *J. Math. Phys.*, 28(9)(1987) 2094–2101.
- [15] W. Hereman and A. Nuseir, Symbolic methods to construct exact solutions of nonlinear partial differential equations, *Mathematics and Computers in Simulation*, 43 (1997), 13–27.
- [16] A.M.Wazwaz, *Partial Differential Equations and Solitary Waves Theorem*, Springer and HEP, Berlin, (2009).
- [17] A.M.Wazwaz, Multiple soliton solutions for coupled KdV and coupled KP Systems, *Canadian Journal of Physics*, 87( 12) (2010) 1227–1232 .
- [18] A.M.Wazwaz, Multiple soliton solutions for the (2+1)-dimensional asymmetric Nizhanik-Novikov-Veselov equation, *Non-linear Analysis Series A: Theory, Methods & Applications* , 72 (2010) 1314–1318.
- [19] A.M.Wazwaz, The (2+1) and (3+1)-dimensional CBS equation: multiple soliton solutions and multiple singular soliton solutions, *Multiple* , *Zeitschrift fur Naturforschung A (ZNA)*, 65a (2010) 173-181.
- [20] A.M.Wazwaz, Multiple-front solutions for the Burgers equation and the coupled Burgers equations, *Appl. Math. Comput.*, 190 (2007) 1198-1206.

- [21] A.M.Wazwaz, New solitons and kink solutions for the Gardner equation, *Communications in Nonlinear Science and Numerical Simulation*, 12(8) (2007), 1395–1404.
- [22] A.M.Wazwaz, Multiple-soliton solutions for the Boussinesq equation, *Appl. Math. Comput.*, 192 (2007) 479-486.
- [23] A. M. Wazwaz, The Hirota's direct method and the tanh-coth method for multiple-soliton solutions of the Sawada-Kotera-Ito seventh-order equation, *Appl. Math. Comput.*, 199(1) (2008) 133–138.
- [24] A.M.Wazwaz, Multiple-front solutions for the Burgers-Kadomtsev-Petvisahvili equation, *Appl. Math. Comput.*, 200 (2008) 437–443.
- [25] A.M.Wazwaz, Multiple-soliton solutions for the Lax-Kadomtsev-Petvisahvili (Lax-KP) equation, *Appl. Math. Comput.*, 201(1/2)(2008) 168–174.
- [26] A.M.Wazwaz, The Hirota's direct method for multiple-soliton solutions for three model equations of shallow water waves, *Appl. Math. Comput.*, 201(1/2) (2008) 489–503.
- [27] A.M.Wazwaz, Multiple-soliton solutions of two extended model equations for shallow water waves, *Appl. Math. Comput.*, 201(1/2) (2008) 790–799.
- [28] A.M.Wazwaz, Single and multiple-soliton solutions for the (2+1)-dimensional KdV equation, *Appl. Math. Comput.*, 204(1) (2008) 20–26.
- [29] A.M.Wazwaz, Solitons and singular solitons for the Gardner-KP equation, *Appl. Math. Comput.*, 204(1) (2008) 162–169.
- [30] A.M.Wazwaz, Solitary wave solutions of the generalized shallow water wave (GSWW) equation by Hirota's method, tanh-coth method and Exp-function method, *Appl. Math. Comput.*, 202 (2008) 275–286.
- [31] A.M.Wazwaz, Multiple kink solutions and multiple singular kink solutions for (2+1)-dimensional nonlinear models generated by the Jaulent-Miodek hierarchy, *Phys. Lett. A*, 373 (2009) 1844–1846.

### **Author information**

Abdul-Majid Wazwaz, Department of Mathematics, Saint Xavier University, Chicago, IL 60655, USA.  
E-mail: wazwaz@sxu.edu

Received: June 10 , 2011

Accepted: August 12, 2011