

A COMPREHENSIVE STUDY OF COMPLETE GENERALIZED NEW MOCK THETA FUNCTIONS

Swayamprabha Tiwari and Dr. Sameena Saba

Communicated by Ayman Badawi

MSC 2010 Classifications: Primary 33C20; Secondary 33C65.

Keywords and phrases: Mock Theta Functions, Bilateral Series, Continued Fractions.

I am thankful to my guide in my research work . My MCN is IU/R&D/2023-MCN0002042.

Corresponding Author: Swayamprabha Tiwari

Abstract The generalization of new mock theta functions of Andrew's and Bringmann-et-al are given. Further we have given the expansion of these bilateral generalized new mock theta functions as ${}_2\phi_1$ series by Slater's transformation. After that we have given the continued fraction representation of these generalized mock theta functions.

1 Introduction

In his last letter to G. H. Hardy [15], S. Ramanujan listed seventeen mock theta functions of order three, five, six and seven. According to Ramanujan a mock theta function is a function $f(q), |q| < 1$, satisfying the following two conditions:

- (0) For every root of unity ξ , there is a θ -function $\theta_\xi(q)$ such that the difference $f(q) - \theta_\xi(q)$ is bounded as $q \rightarrow \xi$ radially.
- (1) There is no single θ -function which works for all ξ , i.e., for every θ -function $\theta(q)$ there is some root of unity ξ for which $f(q) - \theta(q)$ is unbounded as $q \rightarrow \xi$ radially.

After studying mock theta functions, Andrews [14] and Bringmann, Hikami and Lovejoy [5] generated some new mock theta functions. In which four functions of Andrews and two of Bringmann, Hikami and Lovejoy found interesting. We have made comprehensive study of these mock theta functions [2-4].we also found [11-13] is useful in our results.

The four mock theta functions of Andrews [14]:

$$\bar{\psi}_0(q) = \sum_{n=0}^{\infty} \frac{q^{2n^2}}{(-q; q)_{2n}} \quad (1.1)$$

$$\bar{\psi}_1(q) = \sum_{n=0}^{\infty} \frac{q^{2n^2+2n}}{(-q; q)_{2n+1}} \quad (1.2)$$

$$\bar{\psi}_2(q) = \sum_{n=0}^{\infty} \frac{q^{2n^2+2n} (q; q^2)_n}{(q^2; q^2)_n (-q; q)_{2n}} \quad (1.3)$$

$$\bar{\psi}_3(q) = \sum_{n=0}^{\infty} \frac{q^{n^2} (-q; q)_n^2}{(q; q)_{2n}}. \quad (1.4)$$

The two mock theta functions of Bringmann, Hikami and Lovejoy[5] :

$$\bar{\phi}_0(q) = \sum_{n=0}^{\infty} q^n (-q; q)_{2n+1} \quad (1.5)$$

$$\bar{\phi}_1(q) = \sum_{n=0}^{\infty} q^n (-q; q)_{2n}. \quad (1.6)$$

If we extend the definition of the defining series of a mock theta function to negative n , the series will become bilateral and we have Bilateral Mock Theta Functions. Watson called them “Complete”.

2 Notations

In this paper we have use the following notations:

$$\begin{aligned} (a; q^k)_n &= \prod_{m=1}^n \left(1 - aq^{k(m-1)}\right), |q^k| < 1, n, \text{ non-negative integer} \\ (a; q^k)_0 &= 1 (a; q^k)_{\infty} = \prod_{m=1}^{\infty} \left(1 - aq^{k(m-1)}\right), \\ (a_1, a_2, a_3, \dots, a_m; q^k)_n &= (a_1; q^k)_n (a_2; q^k)_n \dots (a_m; q^k)_n \\ &\times {}_A\phi_{A-1} \left[\begin{matrix} a_1, a_2, \dots, a_A; b_1, b_2, \dots, b_{A-1}; q_1, z \end{matrix} \right] \\ &= \sum_{n=0}^{\infty} \frac{(a_1; q_1)_n \dots (a_A; q_1)_n z^n}{(b_1; q_1)_n \dots (b_{A-1}; q_1)_n (q_1; q_1)_n}, |z| < 1 \end{aligned}$$

3 Generalized New mock theta functions

We have given the Generalization of new mock theta functions (1.1) - (1.6) as follows:

$$\bar{\psi}_0(t, \alpha, z; q) = \frac{1}{(t)_{\infty}} \sum_{n=0}^{\infty} \frac{(t)_n q^{2n^2-3n+n\alpha} z^{2n}}{\left(\frac{-z^2}{q}; q\right)_{2n}} \quad (3.1)$$

$$\bar{\psi}_1(t, \alpha, z; q) = \frac{1}{(t)_{\infty}} \sum_{n=0}^{\infty} \frac{(t)_n q^{2n^2-n+n\alpha} z^{2n}}{\left(\frac{-z^2}{q}; q\right)_{2n+1}} \quad (3.2)$$

$$\bar{\psi}_2(t, \alpha, z; q) = \frac{1}{(t)_{\infty}} \sum_{n=0}^{\infty} \frac{(t)_n q^{2n^2+2n-2n\alpha} z^{2n} (z; q^2)_n}{(z^2; q^2)_n \left(\frac{-z^2}{q}; q\right)_{2n}} \quad (3.3)$$

$$\bar{\psi}_3(t, \alpha, z; q) = \frac{1}{(t)_{\infty}} \sum_{n=0}^{\infty} \frac{(t)_n q^{n^2-n+n\alpha} z^{2n} (-z; q)_n^2}{\left(\frac{z^2}{q}; q\right)_{2n}} \quad (3.4)$$

$$\bar{\phi}_0(t, \alpha, z; q) = \frac{1}{(t)_{\infty}} \sum_{n=0}^{\infty} (t)_n q^{n-2n\alpha} z^{2n} \left(\frac{-z^2}{q}; q\right)_{2n+1} \quad (3.5)$$

$$\bar{\phi}_1(t, \alpha, z; q) = \frac{1}{(t)_{\infty}} \sum_{n=0}^{\infty} (t)_n q^{n-2n\alpha} z^{2n} \left(\frac{-z^2}{q}; q\right)_{2n}. \quad (3.6)$$

Now the complete form of new mock theta functions are

$$\bar{\psi}_{0,c}(t, \alpha, z; q) = \frac{1}{(t)_\infty} \sum_{n=-\infty}^{\infty} \frac{(t)_n q^{2n^2-3n+n\alpha} z^{2n}}{\left(\frac{-z^2}{q}; q\right)_{2n}} \quad (3.7)$$

$$\bar{\psi}_{1,c}(t, \alpha, z; q) = \frac{1}{(t)_\infty} \sum_{n=-\infty}^{\infty} \frac{(t)_n q^{2n^2-n+n\alpha} z^{2n}}{\left(\frac{-z^2}{q}; q\right)_{2n+1}} \quad (3.8)$$

$$\bar{\psi}_{2,c}(t, \alpha, z; q) = \frac{1}{(t)_\infty} \sum_{n=-\infty}^{\infty} \frac{(t)_n q^{2n^2+2n-2n\alpha} z^{2n}}{(z^2; q)_n \left(\frac{-z^2}{q}; q\right)_{2n}} \quad (3.9)$$

$$\bar{\psi}_{3,c}(t, \alpha, z; q) = \frac{1}{(t)_\infty} \sum_{n=-\infty}^{\infty} \frac{(t)_n q^{n^2-n+n\alpha} z^{2n} (-z; q)_n^2}{\left(\frac{z^2}{q}; q\right)_{2n}} \quad (3.10)$$

$$\bar{\phi}_{0,c}(t, \alpha, z; q) = \frac{1}{(t)_\infty} \sum_{n=-\infty}^{\infty} (t)_n q^{n-2n\alpha} z^{2n} \left(\frac{-z^2}{q}; q\right)_{2n+1} \quad (3.11)$$

$$\bar{\phi}_{1,c}(t, \alpha, z; q) = \frac{1}{(t)_\infty} \sum_{n=-\infty}^{\infty} (t)_n q^{n-2n\alpha} z^{2n} \left(\frac{-z^2}{q}; q\right)_{2n} \quad (3.12)$$

4 Expansion of Bilateral Generalized New Mock Theta Functions

In the expansion of Slater [8, (5.4.3), p.129] taking $r = 2$, we get:

$$\begin{aligned} & \frac{\left(b_1, b_2, \frac{q}{a_1}, \frac{q}{a_2}, dz, \frac{q}{dz}; q\right)_\infty}{\left(c_1, c_2, \frac{q}{c_1}, \frac{q}{c_2}; q\right)_\infty} {}_2\Psi_2 \left[\begin{matrix} a_1, & a_2 \\ b_1, & b_2 \end{matrix}; q, z \right] = \\ & \frac{q}{c_1} \frac{\left(\frac{c_1}{a_1}, \frac{c_1}{a_2}, \frac{qb_1}{c_1}, \frac{qb_2}{c_1}, \frac{dc_1z}{q}, \frac{q^2}{dc_1z}; q\right)_\infty} {\left(c_1, \frac{q}{c_1}, \frac{c_1}{c_2}, \frac{qc_2}{c_1}; q\right)_\infty} {}_2\Psi_2 \left[\begin{matrix} \frac{qa_1}{c_1}, & \frac{qa_2}{c_1} \\ \frac{qb_1}{c_1}, & \frac{qb_2}{c_1} \end{matrix}; q, z \right] + \text{idem } (c_1; c_2). \end{aligned} \quad (4.1)$$

Where $d = \frac{a_1 a_2}{c_1 c_2}$, $\left| \frac{b_1 b_2}{a_1 a_2} \right| < |z| < 1$

Expansion for equation (3.7) - (3.12) bilateral new generalized mock theta functions are given as follows:

(i) Letting $t \rightarrow 0, q \rightarrow q^2, a_1 a_2 \rightarrow \infty, b_1 = \frac{-z^2}{q}, b_2 = -z^2, d = \frac{a_1 a_2}{c_1 c_2}, z \rightarrow \frac{z^2 q^{\alpha-1}}{a_1 a_2}$ in we get:

$$\begin{aligned} & \frac{\left(\frac{-z^2}{q}, -z^2, \frac{z^2 q^{\alpha-1}}{c_1 c_2}, \frac{c_1 c_2}{z^2 q^{\alpha-3}}; q^2\right)_\infty}{\left(c_1, c_2, \frac{q^2}{c_1}, \frac{q^2}{c_2}; q^2\right)_\infty} \cdot \bar{\psi}_{0,c}(0, \alpha, z; q) \\ &= \frac{q^2}{c_1} \frac{\left(\frac{-z^2 q}{c_1}, \frac{-z^2 q^2}{c_1}, \frac{z^2 q^{\alpha-3}}{c_1 c_2}, \frac{c_1 q^{5-\alpha}}{z^2}; q^2\right)_\infty}{\left(c_1, \frac{q^2}{c_1}, \frac{c_1}{c_2}, \frac{q^2 c_2}{c_1}; q^2\right)_\infty} \\ & \times \frac{1}{c_1^{2n}} \sum_{n=-\infty}^{\infty} \frac{q^{2n^2+n+n\alpha} z^{2n}}{\left(\frac{-z^2 q}{c_1}; q^2\right)_n \left(\frac{-z^2 q^2}{c_1}; q^2\right)_n} + \text{idem } (c_1, c_2) \end{aligned} \quad (4.2)$$

(ii) Letting $t \rightarrow 0, q \rightarrow q^2, a_1 a_2 \rightarrow \infty, b_1 = -z^2, b_2 = -z^2 q, d = \frac{a_1 a_2}{c_1 c_2}, z \rightarrow \frac{z^2 q^{\alpha+1}}{a_1 a_2}$ in (4.1) we get:

$$\begin{aligned}
& \frac{\left(1 + \frac{z^2}{q}\right)(-z^2, -z^2q, \frac{z^2q^{\alpha+1}}{c_1c_2}, \frac{c_1c_2}{z^2q^{\alpha-1}}; q^2)_\infty}{\left(c_1, c_2, \frac{q^2}{c_1}, \frac{q^2}{c_2}; q^2\right)_\infty} \cdot \bar{\psi}_{1,c}(0, \alpha, z; q) \\
&= \frac{q^2}{c_1} \frac{\left(\frac{-z^2q^2}{c_1}, \frac{-z^2q^3}{c_1}, \frac{z^2q^{\alpha-1}}{c_2}, \frac{c_2}{z^2q^{\alpha-3}}; q^2\right)_\infty}{\left(c_1, \frac{q^2}{c_1}, \frac{c_1}{c_2}, \frac{q^2c_2}{c_1}; q^2\right)_\infty} \\
&\quad \times \frac{1}{c_1^{2n}} \sum_{n=-\infty}^{\infty} \frac{q^{2n^2+3n+n\alpha} z^{2n}}{\left(\frac{-z^2q^2}{c_1}; q^2\right)_n \left(\frac{-z^2q^3}{c_1}; q^2\right)_n} + \text{idem } (c_1, c_2) \quad (4.3)
\end{aligned}$$

(iii) Letting $t \rightarrow 0, q \rightarrow q^2, a_1 = -z^2, a_2 = -z^2q, b_1 = b_2 = 0, d = \frac{a_1a_2}{c_1c_2}, z \rightarrow z^2q^{1-2\alpha}$ in (4.1) we get:

$$\begin{aligned}
& \frac{\left(\frac{-q^2}{z^2}, \frac{-q}{z^2}, \frac{z^6q^{2(1-\alpha)}}{c_1c_2}, \frac{c_1c_2}{z^4q^{-2\alpha}}; q^2\right)_\infty}{\left(1 + \frac{z^2}{q}\right)(c_1, c_2, \frac{q^2}{c_1}, \frac{q^2}{c_2}; q^2)_\infty} \cdot \bar{\phi}_{0,c}(0, \alpha, z; q) \\
&= \frac{q^2}{c_1} \frac{\left(\frac{-c_1}{z^2}, \frac{-c_1}{z^2q}, \frac{z^6}{q^{2\alpha}c_2}, \frac{c_2q^{2+2\alpha}}{z^6}; q^2\right)_\infty}{\left(c_1, \frac{q^2}{c_1}, \frac{c_1}{c_2}, \frac{q^2c_2}{c_1}; q^2\right)_\infty} \\
&\quad \times \sum_{n=-\infty}^{\infty} q^{n-2n\alpha} \left(\frac{-z^2q^2}{c_1}; q^2\right)_n \left(\frac{-z^2q^3}{c_1}; q^2\right)_n z^{2n} + \text{idem } (c_1, c_2) \quad (4.4)
\end{aligned}$$

(iv) Letting $t \rightarrow 0, q \rightarrow q^2, a_1 = \frac{-z^2}{q}, a_2 = -z^2, b_1 = b_2 = 0, d = \frac{a_1a_2}{c_1c_2}, z \rightarrow z^2q^{1-2\alpha}$ in we get:

$$\begin{aligned}
& \frac{\left(\frac{-q^3}{z}, \frac{-q^2}{z^2}, \frac{z^6}{c_1c_2q^{2\alpha}}, \frac{c_1c_2q^{2+2\alpha}}{z^6}; q^2\right)_\infty}{\left(c_1, c_2, \frac{q^2}{c_1}, \frac{q^2}{c_2}; q^2\right)_\infty} \cdot \bar{\phi}_{1,c}(0, \alpha, z; q) \\
&= \frac{q^2}{c_1} \frac{\left(\frac{-c_1}{z^2}, \frac{-qc_1}{z^2} \frac{z^6q^{-2-2\alpha}}{c_2}, \frac{c_2q^{4+2\alpha}}{z^6}; q^2\right)_\infty}{\left(c_1, \frac{q^2}{c_1}, \frac{c_1}{c_2}, \frac{q^2c_2}{c_1}; q^2\right)_\infty} \\
&\quad \times \sum_{n=-\infty}^{\infty} q^{n-2n\alpha} \left(\frac{-z^2q}{c_1}; q^2\right)_n \left(\frac{-z^2q^2}{c_1}; q^2\right)_n z^{2n} + \text{idem } (c_1, c_2) \quad (4.5)
\end{aligned}$$

5 Expansion for $\bar{\psi}_{0,c}(t, \alpha, z; q)$ and $\bar{\psi}_{1,c}(t, \alpha, z; q)$ as ${}_2\phi_1$ Series

Taking $j = 1, 2$ in [8, (5.4.5).p.130] we get:

$$\begin{aligned}
& \frac{\left(\frac{q}{a_1}, \frac{q}{a_2}, dz, \frac{q}{dz}; q\right)_\infty}{{\left(\frac{q}{b_1}, \frac{q}{b_2}; q\right)}_\infty} {}_2\psi_2 \left[\begin{matrix} a_1, & a_2 \\ b_1, & b_2 \end{matrix}; q, z \right] = \\
& \quad \frac{q}{b_1} \frac{\left(q, \frac{b_1}{a_1}, \frac{b_1}{a_2}, \frac{db_1z}{q}, \frac{q^2}{db_1}; q\right)_\infty}{{\left(b_1, \frac{q}{b_1}, \frac{b_1}{b_2}; q\right)}_\infty} {}_2\phi_1 \left[\begin{matrix} \frac{qa_1}{b_1}, & \frac{qa_2}{b_1} \\ \frac{qb_2}{b_1} \end{matrix}; q, z \right] + \text{idem } (b_1; b_2) \quad (5.1)
\end{aligned}$$

Where $d = \frac{a_1a_2}{b_1b_2}$. By using (5.1) we have given expansion of bilateral new generalized mock theta functions (3.7) – (3.8) as ${}_2\phi_1$ series:

(i) Letting $q \rightarrow q^2, a_1a_2 \rightarrow \infty, t \rightarrow 0, b_1 = \frac{-z^2}{q}, b_2 = -z^2$ and $z \rightarrow \frac{z^2q^{\alpha-1}}{a_1a_2}$ in (5.1) we get:

$$\begin{aligned}\bar{\psi}_{0,c}(0, \alpha, z; q) &= \frac{-q^3}{z^2} \frac{\left(q^2, -q^{\alpha-3}, -q^{3-\alpha}, \frac{-q^2}{z^2}; q^2\right)_\infty}{\left(\frac{-z^2}{q}, \frac{1}{q}, \frac{q^\alpha}{z^2}, z^2 q^{2-\alpha}; q^2\right)_\infty} \\ &\quad \times \sum_{n=0}^{\infty} \frac{q^{2n^2+3n+n\alpha} z^{-2n}}{(q^3; q^2)_n (q^2; q^2)_n} + \text{idem } (b_1, b_2) \quad (5.2)\end{aligned}$$

(ii) Letting $q \rightarrow q^2, a_1 a_2 \rightarrow \infty, t \rightarrow 0, b_1 = -z^2, b_2 = -z^2 q$ and $z \rightarrow \frac{z^2 q^{\alpha+1}}{a_1 a_2}$ in (5.1) we get :

$$\begin{aligned}\bar{\psi}_{1,c}(0, \alpha, z; q) &= \frac{-q^2}{z^2} \frac{\left(q^2, -q^{\alpha-2}, -q^{4-\alpha}, \frac{-q^2}{z^2}; q^2\right)_\infty}{\left(-z^2, \frac{1}{q}, \frac{q^\alpha}{z^2}, z^2 q^{2-\alpha}; q^2\right)_\infty} \\ &\quad \times \sum_{n=0}^{\infty} \frac{q^{2n^2+3n+n\alpha} z^{-2n}}{(q^3; q^2)_n (q^2; q^2)_n} + \text{idem } (b_1, b_2) \quad (5.3)\end{aligned}$$

6 Expansion of $\bar{\phi}_{0,c}(t, \alpha, z; q)$ and $\bar{\phi}_{1,c}(t, \alpha, z; q)$ as ${}_2\phi_1$ Series

For getting the expansion of a bilateral new generalized mock theta functions (3.11) (3.12) as a ${}_2\phi_1$ series, we take $c_j = qa_j, j = 1, 2$ in Slater's expansion formula [8, (5.4.4), p.130] we get :

$$\begin{aligned}&\frac{\left(b_1, b_2, \frac{q}{a_1}, \frac{q}{a_2}, z, \frac{q}{z}; q\right)_\infty} {\left(qa_1, qa_2, \frac{1}{a_1}, \frac{1}{a_2}; q\right)_\infty} {}_2\psi_2 \left[\begin{matrix} a_1, & a_2 \\ b_1, & b_2 \end{matrix}; q, z \right] \\ &= \frac{a_1 \left(q, \frac{qa_1}{a_2}, \frac{b_1}{a_1}, \frac{b_2}{a_1}, a_1 z, \frac{q}{a_1 z}; q\right)_\infty} {\left(qa_1, \frac{1}{a_1}, \frac{a_1}{a_2}, \frac{qa_2}{a_1}; q\right)_\infty} {}_2\phi_1 \left[\begin{matrix} \frac{qa_1}{b_1}, \frac{qa_1}{b_2} \\ \frac{qa_1}{a_2}; q \frac{b_1 b_2}{a_1 a_2 z} \end{matrix} \right] + \text{idem } (a_1, a_2), \quad (6.1)\end{aligned}$$

where $\left| \frac{b_1 b_2}{a_1 a_2} \right| < |z| < 1$

(i) Letting $q \rightarrow q^2, b_1 = b_2 = 0, a_1 = -z^2, a_2 = -z^2 q, d = \frac{a_1 a_2}{c_1 c_2}, z \rightarrow z^2 q^{1-2\alpha}$ in (6.1), we have

$$\begin{aligned}\bar{\phi}_{0,c}(0, \alpha, z; q) &= \frac{-z^2(1-q) \left(q^2, -z^4 q^{1-2\alpha}, \frac{-q^{1+2\alpha}}{z^4}, -z^2 q^3, -z^2 q^2, \frac{-1}{z^2 q}, \frac{-1}{z^2}; q^2\right)_\infty}{\left(\frac{1}{q}, \frac{-q^2}{z^2}, \frac{-q}{z}, z^2 q^{1-2\alpha}, \frac{q^{1+2\alpha}}{z^2}; q^2\right)_\infty} \\ &\quad \times \sum_{n=0}^{\infty} \frac{q^{2n^2+2n\alpha} z^{-2n}}{(q; q^2)_n (q^2; q^2)_n} + \text{idem } (a_1, a_2) \quad (6.2)\end{aligned}$$

(ii) Letting $q \rightarrow q^2, b_1 = b_2 = 0, a_1 = \frac{-z^2}{q}, a_2 = -z^2, d = \frac{a_1 a_2}{c_1 c_2}, z \rightarrow z^2 q^{1-2\alpha}$ in (6.1), we have

$$\begin{aligned}\bar{\phi}_{1,c}(0, \alpha, z; q) &= \frac{-z^2}{q} (1-q) \left(1 + \frac{q}{z^2}\right) \left(1 + \frac{1}{z^2}\right) \frac{\left(q^2, -z^4 q^{-2\alpha}, \frac{-q^{2+2\alpha}}{z^4}, -z^2 q^2; q^2\right)_\infty}{\left(\frac{1}{q}, \frac{-q}{z^2}, z^2 q^{1-2\alpha}, \frac{q^{1+2\alpha}}{z^2}; q^2\right)_\infty} \\ &\quad \times \sum_{n=0}^{\infty} \frac{q^{2n^2+2n} z^{-2n}}{(q; q^2)_n (q^2; q^2)_n} + \text{idem } (a_1, a_2) \quad (6.3)\end{aligned}$$

7 Bilateral Generalized New Mock Theta Functions as continued fractions

By using the continued fraction [14, (3.79), p.82] given by

$$\frac{\sum_{n=0}^{\infty} \frac{q^{n^2} \lambda^n}{(q)_n (-\beta)_n}}{\sum_{n=0}^{\infty} \frac{q^{n^2+n} \lambda^n}{(q)_n (-\beta)_n}} = 1 + \frac{\lambda q + \beta}{1 - \beta} \cdot \frac{\lambda q^2 + \beta}{1 - \beta} \dots, \quad (7.1)$$

We have given the continued fraction representation for $\bar{\psi}_{0,c}(t, \alpha, z; q)$, $\bar{\psi}_{1,c}(t, \alpha, z; q)$.

(i) Representation of $\bar{\psi}_{0,c}(t, \alpha, z; q)$ as a continued fraction.

Now (5.2) gives

$$\begin{aligned} \bar{\psi}_{0,c}(0, \alpha, z; q) &= \frac{-q^3}{z^2} \frac{\left(q^2, -q^{\alpha-3}, -q^{3-\alpha}, \frac{-q^2}{z^2}; q^2 \right)_\infty}{\left(\frac{-z^2}{q}, \frac{1}{q}, \frac{q^\alpha}{z^2}, z^2 q^{2-\alpha}; q^2 \right)_\infty} \times \sum_{n=0}^{\infty} \frac{q^{2n^2+3n+n\alpha} z^{-2n}}{(q^3; q^2)_n (q^2; q^2)_n} \\ &\quad + \frac{-q^2}{z^2} \frac{\left(q^2, -q^{\alpha-2}, -q^{2-\alpha}, \frac{-q^3}{z^2}; q^2 \right)_\infty}{\left(-z^2, z^2 q^{\alpha-2}, \frac{q^\alpha}{z^2}, q; q^2 \right)_\infty} \times \sum_{n=0}^{\infty} \frac{q^{2n^2+n+n\alpha} z^{-2n}}{(q; q^2)_n (q^2; q^2)_n} \\ &= S \sum_{n=0}^{\infty} \frac{q^{2n^2+3n+n\alpha} z^{-2n}}{(q^3; q^2)_n (q^2; q^2)_n} + T \sum_{n=0}^{\infty} \frac{q^{2n^2+n+n\alpha} z^{-2n}}{(q; q^2)_n (q^2; q^2)_n} \end{aligned} \quad (7.2)$$

Where

$$S = \frac{-q^3}{z^2} \frac{\left(q^2, -q^{\alpha-3}, -q^{3-\alpha}, \frac{-q^2}{z^2}; q^2 \right)_\infty}{\left(\frac{-z^2}{q}, \frac{1}{q}, \frac{q^\alpha}{z^2}, z^2 q^{2-\alpha}; q^2 \right)_\infty}$$

and

$$T = \frac{-q^2}{z^2} \frac{\left(q^2, -q^{\alpha-2}, -q^{2-\alpha}, \frac{-q^3}{z^2}; q^2 \right)_\infty}{\left(-z^2, z^2 q^{\alpha-2}, \frac{q^\alpha}{z^2}, q; q^2 \right)_\infty}$$

Dividing the equation (7.2) by the first series on the right hand side of (7.2) we get;

$$\frac{\bar{\psi}_{0,c}(0, \alpha, z; q)}{\sum_{n=0}^{\infty} \frac{q^{2n^2+3n+n\alpha} z^{-2n}}{(q^3; q^2)_n (q^2; q^2)_n}} = S + T \frac{\sum_{n=0}^{\infty} \frac{q^{2n^2+n+n\alpha} z^{-2n}}{(q; q^2)_n (q^2; q^2)_n}}{\sum_{n=0}^{\infty} \frac{q^{2n^2+3n+n\alpha} z^{-2n}}{(q^3; q^2)_n (q^2; q^2)_n}}. \quad (7.3)$$

Taking $q \rightarrow q^2$, $\lambda \rightarrow z^{-2} q^{1+\alpha}$, $\beta = -q$ in (7.1), and then putting it in the quotient of the summation on the right side of (7.3) to get the continued fraction representation for $\bar{\psi}_{0,c}(t, \alpha, z; q)$ as:

$$\frac{\bar{\psi}_{0,c}(0, \alpha, z; q)}{\sum_{n=0}^{\infty} \frac{q^{2n^2+3n+n\alpha} z^{-2n}}{(q^3; q^2)_n (q^2; q^2)_n}} = S + \frac{T}{(1-q)} \left[1 + \frac{q^{\alpha+3} z^{-2} - q}{(1+q)} \cdot \frac{q^{\alpha+5} z^{-2} - q}{(1+q)} + \dots \right] \quad (7.4)$$

(ii) Representation of $\bar{\psi}_{1,c}(t, \alpha, z; q)$ as a continued fraction similar as earlier one.

$$\begin{aligned}\bar{\psi}_{1,c}(0, \alpha, z; q) &= \frac{-q^2}{z^2} \frac{\left(q^2, -q^{\alpha-2}, -q^{4-\alpha}, \frac{-q^2}{z^2}; q^2\right)_\infty}{\left(-z^2, \frac{1}{q}, \frac{q^\alpha}{z^2}, z^2q^{2-\alpha}; q^2\right)_\infty} \sum_{n=0}^{\infty} \frac{q^{2n^2+3n+n\alpha} z^{-2n}}{(q^3; q^2)_n (q^2; q^2)_n} \\ &+ \frac{-q}{z^2} \frac{\left(q^2, -q^{\alpha-1}, -q^{3-\alpha}, \frac{-q^2}{z^2}; q^2\right)_\infty}{\left(-z^2q, q, z^2q^{2-\alpha}, \frac{q^\alpha}{z^2}; q^2\right)_\infty} \sum_{n=0}^{\infty} \frac{q^{2n^2+n+n\alpha} z^{-2n}}{(q; q^2)_n (q^2; q^2)_n} \\ &= S_1 \sum_{n=0}^{\infty} \frac{q^{2n^2+3n+n\alpha} z^{-2n}}{(q^3; q^2)_n (q^2; q^2)_n} + T_1 \sum_{n=0}^{\infty} \frac{q^{2n^2+n+n\alpha} z^{-2n}}{(q; q^2)_n (q^2; q^2)_n}\end{aligned}$$

Where

$$\begin{aligned}S_1 &= \frac{-q^2}{z^2} \frac{\left(q^2, -q^{\alpha-2}, -q^{4-\alpha}, \frac{-q^2}{z^2}; q^2\right)_\infty}{\left(-z^2, \frac{1}{q}, \frac{q^\alpha}{z^2}, z^2q^{2-\alpha}; q^2\right)_\infty} \\ T_1 &= \frac{-q}{z^2} \frac{\left(q^2, -q^{\alpha-1}, -q^{3-\alpha}, \frac{-q^2}{z^2}; q^2\right)_\infty}{\left(-z^2q, q, z^2q^{2-\alpha}, \frac{q^\alpha}{z^2}; q^2\right)_\infty}\end{aligned}$$

Hence we have

$$\frac{\bar{\psi}_{1,c}(0, \alpha, z; q)}{\sum_{n=0}^{\infty} \frac{q^{2n^2+3n+n\alpha} z^{-2n}}{(q^3; q^2)_n (q^2; q^2)_n}} = S_1 + \frac{T_1}{(1-q)} \left[1 + \frac{q^{3+\alpha} z^{-2} - q}{1+q} \cdot \frac{q^{5+\alpha} z^{-2} - q}{1+q} \right]. \quad (7.5)$$

8 Conclusion remarks

The study of bilateral generalized new mock theta functions is interesting as it gives the expansion by Slater and also expansion for these generalized new mock theta functions as ${}_2\phi_1$ Series. After that we have given continued fractions for generalized new mock theta functions.

References

- [1] B.Gordon and R.J. McIntosh, Some eighth order mock theta functions, J. London Math. Soc. 2000,62 (2): 321-335.
- [2] B. Srivastava, A generalization and study of four new mock theta functions, American Journal of Computational and Applied Mathematics 2012, 2(2):58-63.
- [3] B. Srivastava, A Study of Bilateral New Mock Theta Functions, American Journal of Mathematics and Statistics 2012, 2(4):64-69.
- [4] B. Srivastava, Partial new mock theta functions, Applied Mathematics 2012, 2(4):146- 151.
- [5] E. Mortess, On three third order mock theta functions and Hecke-type double sums (pre –print).
- [6] G. E. Andrews, q-orthogonal polynomials, Rogers- Ramanujan identities, and mock theta functions, preprint.
- [7] G. E. Andrews, D. Hickerson, Ramanujan's "Lost" Notebook VII: The sixth order mock theta functions, Adv. in Math. 89 (1991), 60-105.
- [8] G. Gasper and M. Rahman, Basic Hypergeometric Series, Cambridge University Press, Cambridge (1990).
- [9] Mohd. Ahmad , S. Faruqi , Some Bilateral Mock Theta Functions and their Lerch Representations, The Aligrah Bulletin of Mathematics, 2015, 34:75-92.
- [10] Mohd. Ahmad , S. Haq , A. H. Khan , Bilateral Mock Theta Functions and Further Properties, Electronic Journal of mathematical analysis and applications 2019,7(2):216-229.
- [11] M.I.Qureshi,M.Shadab and M.S.Baboo,Evaluation of some novel integrals involving legendre function of second kind using hypergeometric approach, Palestine Journal of Mathematics 2017,6(1):68-75.
- [12] P.K.Ray and Juli Sahu,Generating function for certain balancing and lucas-balancing numbers, Palestine Journal of Mathematics 2016,5(2):122-129.

- [13] P.Murugan and S.N.Fathima,Some congruences for ramanujan's general partition function,Palestine Journal of Mathematics 2022,11(2):38-44.
- [14] R.P. Agarwal, Resonance of Ramanujan's Mathematics III, New Age International, New Delhi, 1996.
- [15] S.Ramanujan, Collected Papers, Cambridge University Press, 1972, reprinted Chelsea, New York, 1962.

Author information

Swayamprabha Tiwari, Department of Mathematics & Statistics, Faculty of Science, Integral University, Lucknow-226026, India.
E-mail: swayam6288@gmail.com

Dr. Sameena Saba, Department of Mathematics & Statistics, Faculty of Science, Integral University, Lucknow-226026, India.
E-mail: sameena9554@gmail.com

Received: 2023-10-05

Accepted: 2024-02-15